

# **Historic, Archive Document**

Do not assume content reflects current  
scientific knowledge, policies, or practices.



a 5011  
A 42 Minervos  
Und 2/11/85 Sch



United States  
Department of  
Agriculture

Forest Service

Rocky Mountain  
Forest and Range  
Experiment Station

Fort Collins,  
Colorado 80526

General Technical  
Report RM-113



# The Vegetation of Theodore Roosevelt National Park, North Dakota: A Habitat Type Classification

Paul L. Hansen, George R. Hoffman, and Ardell J. Bjugstad

U.S. DEPT. OF AGRICULTURE  
APR 3 1985



### **Abstract**

Vegetation of the Theodore Roosevelt National Park, North Dakota, was characterized according to habitat type based on concepts and methods developed by Daubenmire. Ten habitat types were described: seven—three grassland and four wooded types—occurred on upland plateaus, slopes, and relatively narrow benches along contours of the hills above the stream valleys; three—all wooded—occurred in valley bottoms. A key to identify the habitat types and discussion on the validation of the Habitat Type Classification and Species Diversity are provided.

### **Acknowledgements**

This study was done with financial support of grants CX-1200-8-B030 and CX-1200-9-B037 from the University of Wyoming-National Park Service Research Center. Bob Powell, Chief Ranger, Theodore Roosevelt National Park, and his staff cooperated and provided a number of services during the summers of 1978 and 1979 when the field studies were being done there. Theodore Van Bruggen, University of South Dakota, provided expertise in the identification of a number of species. The authors also thank Kenneth L. Diem, Director, University-National Park Service Research Center, for his cooperation and support in the publication of this report.

# **The Vegetation of Theodore Roosevelt National Park, North Dakota: A Habitat Type Classification**

**Paul L. Hansen, Research Assistant  
George R. Hoffman, Professor of Biology  
Department of Biology, University of South Dakota  
and  
Ardell J. Bjugstad, Supervisory Range Scientist  
Rocky Mountain Forest and Range Experiment Station<sup>1</sup>**

<sup>1</sup>*Station headquarters is in Fort Collins, in cooperation with Colorado State University. The Station's Research Work Unit at Rapid City cooperated with the University of South Dakota at Vermillion in analyzing the research.*

## Contents

	Page
STUDY AREA .....	1
Physiography and Soils .....	1
Climate .....	1
ECOLOGIC TERMS AND CONCEPTS .....	2
PREVIOUS VEGETATION STUDIES .....	2
METHODS .....	3
HABITAT TYPE DESCRIPTIONS .....	4
<i>Stipa comata</i> / <i>Carex filifolia</i> Habitat Type .....	4
<i>Agropyron smithii</i> / <i>Carex filifolia</i> Habitat Type .....	5
<i>Andropogon scoparius</i> / <i>Carex filifolia</i> Habitat Type .....	6
<i>Juniperus horizontalis</i> / <i>Andropogon scoparius</i> Habitat Type .....	7
<i>Artemisia tridentata</i> / <i>Agropyron smithii</i> Habitat Type .....	8
<i>Artemisia cana</i> / <i>Agropyron smithii</i> Habitat Type .....	9
<i>Fraxinus pennsylvanica</i> / <i>Symphoricarpos occidentalis</i> Habitat Type .....	11
<i>Fraxinus pennsylvanica</i> / <i>Prunus virginiana</i> Habitat Type .....	12
<i>Populus tremuloides</i> / <i>Betula occidentalis</i> Habitat Type .....	14
<i>Juniperus scopulorum</i> / <i>Oryzopsis micrantha</i> Habitat Type .....	15
<i>Symphoricarpos occidentalis</i> Community .....	15
KEY TO THE HABITAT TYPES .....	17
DISCUSSION .....	18
Validation of the Habitat Type Classification .....	18
Species Diversity .....	18
LITERATURE CITED .....	19
APPENDIX 1. Habitat Type Tables with Stand Data .....	21
APPENDIX 2. Tree Population Analyses .....	32
APPENDIX 3. Soil Analyses .....	34



# The Vegetation of Theodore Roosevelt National Park, North Dakota: A Habitat Type Classification

Paul L. Hansen, George R. Hoffman, and Ardell J. Bjugstad

Increasing utilization of the northern Great Plains is reducing the areas of vegetation that are climax, or near-climax. Therefore, it becomes increasingly important to document in detail the vegetation characteristics of those areas currently free from, or only minimally disturbed by, human activities. Such areas are potentially valuable as reference areas against which to measure the impact of human activities. Theodore Roosevelt National Park (TRNP) in western North Dakota represents such a reference area, which also has a considerable diversity of habitats and vegetation types. This study documents the characteristics of pristine, or near-pristine, vegetation, and describes and delimits the habitat types (Daubenmire 1968) of TRNP. The information may be of interest to both those interested in basic ecological data and descriptions and those directly concerned with management of the vegetation described.

## STUDY AREA

### Physiography and Soils

Theodore Roosevelt National Park (TRNP) was established as a National Park in 1947. It covers 26,578 ha divided between the North and South Units (fig. 1). The area is part of the Missouri River Plateau and is unglaciated, except for a small portion of the North Unit, and even there, glacial deposits are not particularly evident. The substrates are stratified beds of soft shales, sandstones, and lignite of the Tertiary Tongue River Formation of the Fort Union Group (Laird 1950, 1956). These substrates have been severely dissected by the Little Missouri River and its tributaries, resulting in "badlands" topography of gullied ravines and valleys separated by interstream uplands of various dimensions, from large plateaus of many hectares to small buttes of erosion-resistant sandstone or scoria. Some of the buttes are 200 m above the valley floors. Exposed lignite has eroded along some of the steep hills, leaving resistant terraces paralleling the contours of the hills. At lower levels, just above the floodplains of the major streams, terraces of alluvium add another distinctive feature to the landscape. The region is physiographically complex, with various sites for the establishment and maintenance of vegetation types (fig. 2). The soils of TRNP are regosols belonging to the Bainville Series, developed from excessively drained, medium-textured, calcareous parent material (Omodt et al. 1968). A typical and obviously immature soil profile shows an A1 horizon overlying a C horizon. Textures range from loams to clay loams.




Figure 1.—Map of western North Dakota showing location of North and South Units of Theodore Roosevelt National Park.

### Climate

The arid and continental climate of western North Dakota is characterized by long, cold winters and short, warm summers. The mean monthly low for January is  $-11.6^{\circ}\text{C}$ , and the mean monthly high for July is  $22.0^{\circ}\text{C}$ . Most precipitation falls in early summer. June (the wettest month) precipitation ranges from mean 98.3 mm at Belfield to 83.6 mm at Medora (table 1). July means range from 72.6 mm at Belfield to 50.5 mm at Medora, illustrating the large decrease from June to July. The increase from May to June is equally impressive. Of the total precipitation in the region recorded at weather stations, 75% falls from April through September.



Figure 2.—General view of the landscape at TRNP showing the badland topography.

## ECOLOGIC TERMS AND CONCEPTS

The following terms and concepts are consistent with usage proposed by Daubenmire (1968) and used in previous studies of habitat types (Daubenmire and Daubenmire 1968; Daubenmire 1970; Hoffman and Alexander 1976, 1980; Pfister et al. 1977; Mueggler and Stewart 1980). "Climax vegetation" is that which has attained a steady-state with its environment, such that species successfully maintain their population sizes. "Habitat type" is defined as the land area which supports or can support the same primary climax vegetation (Daubenmire 1968). The climax vegetation is referred to as a plant association. As an example, the *Artemisia tridentata*/ *Agropyron spicatum* habitat type of south-central Washington state supports the plant association of the same name, under pristine conditions. Grazing, however, can alter significantly the species composition such that a community of *Artemisia tridentata*/ *Bromus tectorum* can result. If the same plant association were burned, killing *Artemisia tridentata*, and grazed, removing *Agropyron spicatum*, the resulting plant community might become a *Poa secunda*/ *Bromus tectorum* community. The habitat type does not change as long as the habitats retain the capability of supporting the *Artemisia*/ *Agropyron* association (Daubenmire 1970). On normal soils and topography, the climax vegetation is referred to as a "climatic climax." Where slope and soil exert predominant influence on the microclimate, compared to that of normal topography, the climax vegetation may reflect the altered microclimate in both composition and structure. Such self-perpetuating vegetation, if significantly different from that on normal soils and topography, is referred to as a "topographic climax" or an "edaphic climax" or in some cases a "topoedaphic climax," where both slope and soil are important in altering the

climax vegetation. A "zootic climax" occurs where periodic grazing and trampling by domestic animals has altered the vegetation significantly. The altered vegetation maintains its composition and structure as a result of the kind and intensity of the periodic grazing. Seral vegetation is that which has not attained a steady-state with its environment. Ordinarily it follows a major disturbance.

Habitat types are defined to delimit and ecologically characterize land units of similar biotic potential, which makes the concept valuable to both basic and applied science. Habitat types can be useful in predicting rates of tree growth, susceptibility of trees to insects and mistletoe infection, potential for producing browse after fire, depth of soil moisture penetration, drought during the summer, and successional trends after disturbance (Arno and Pfister 1977; Daubenmire 1961, 1972; Hoffman and Alexander 1980, 1983; Layser 1974; Mueggler and Stewart 1980; Pfister 1972). The present study can provide a basis for better understanding the effects of grazing on habitat types in western North Dakota.

## PREVIOUS VEGETATION STUDIES

Previous vegetation studies in western North Dakota did not include attempts to delimit habitat types. Hanson and Whitman (1938) described nine major kinds of grassland vegetation, which they correlated with edaphic and topographic factors. At least three of their types were considered to be seral, and the status of some others was not given. Their *Bouteloua gracilis*/ *Stipa comata*/ *Carex filifolia* vegetation type on upland plateaus and gentle slopes, and the *Artemisia cana*/ *Agropyron smithii* vegetation type of alluvial flats adjacent the Little Missouri River and its tributaries were considered to be stable types. The *Andropogon gerardii*-



Table 1.—Mean precipitation (P, in mm) and temperature (T, in °C) from selected weather stations near Theodore Roosevelt National Park

Month	Location							
	Beach		Belfield		Medora 3NNE		Watford City	
	P	T	P	T	P	T	P	T
Jan.	12.9	−11.1	15.7	−11.8	11.4	−11.9	14.5	−11.4
Feb.	10.4	−8.7	17.0	−8.4	12.7	−7.8	12.7	−9.4
Mar.	15.7	−2.9	22.1	−4.2	16.3	−3.3	19.8	−3.4
Apr.	30.5	5.5	32.5	5.1	25.9	5.7	34.0	6.1
May	55.4	11.7	55.9	11.9	44.2	12.6	48.0	12.9
June	90.2	16.8	98.3	16.3	83.6	17.2	86.9	17.3
July	44.2	20.9	72.6	20.1	50.5	21.2	52.6	21.8
Aug.	40.6	19.7	61.0	19.8	37.1	20.6	38.6	20.6
Sept.	31.5	13.8	30.7	13.6	30.7	14.1	33.3	14.4
Oct.	23.1	7.1	25.9	7.4	20.3	7.4	23.9	7.9
Nov.	12.7	−0.8	19.0	−1.6	14.2	−0.9	14.5	−1.7
Dec.	10.2	−7.2	8.6	−6.7	6.9	−6.9	13.2	−7.5
Annual <sup>1</sup>	377.4	5.4	459.3	5.1	353.8	5.7	392.0	5.7

<sup>1</sup> Mean total annual precipitation.

dominated type, much more restricted in distribution, also might have been stable (Hanson and Whitman 1938). Quinnild and Cosby (1958) found three apparently stable vegetation types on two ungrazed mesas in western North Dakota. The types were dominated by *Agropyron smithii*, *A. dasystachyum*, and *Stipa comata*, respectively. Larson and Whitman (1942) reported a vegetation type dominated by *Agropyron smithii*, *Bouteloua gracilis*, *Carex eleocharis*, and *C. filifolia* on an ungrazed mesa, in the badlands of South Dakota. From the badlands of southeastern Montana, Brown (1971) reported seven vegetation types dominated by the following: *Sarcobatus vermiculatus*, *Atriplex confertifolia*, *Artemisia tridentata*, *Artemisia tridentata*/*Atriplex confertifolia*-*Agropyron spicatum*, *Artemisia tridentata*/*Agropyron spicatum*, *Rhus trilobata*/*Agropyron spicatum*, *Juniperus scopulorum*/*Agropyron spicatum*, *Pinus ponderosa*/*Juniperus scopulorum*. The distribution of most of these vegetation types appeared to be controlled by specific soil and/or topographic characteristics. Ralston (1960) concluded that the lower temperatures and higher moisture levels of north-facing exposures at TRNP were primarily responsible for the success of *Juniperus scopulorum*-dominated vegetation there. Nelson (1961) described two additional forest types at TRNP, including a *Populus deltoides* vegetation type along streams, and a *Fraxinus pennsylvanica* vegetation type along streams and in upland ravines.

## METHODS

The approach used follows that of Daubenmire and Daubenmire (1968) and Daubenmire (1970). Preliminary work at TRNP was done during the summer of 1978, by traveling extensively throughout the North and South Units, collecting plants and noting possible study sites. Stands of vegetation were sought that were nearly mature, most homogeneous, and least disturbed by fire, trails, past farming, or grazing. Homogeneity was an

important criterion and was determined subjectively. Obvious discontinuities and sharp breaks in the microtopography of any stand, and locally disturbed areas, such as animal trails, were avoided. Areas in stands were chosen which appeared to have a representative expression of all vegetation layers present. From the 1978 field work, a tentative list was made of the habitat types to be substantiated or altered as a result of intensive sampling in 1979.

During the summer of 1979, 70 stands were intensively sampled. At each stand sampled, a rectangular plot 15 x 25 m was set up. If the plot was located on a slope, it was oriented with the long axis parallel to the contour to maximize the chance of remaining within the same soil type. In woodland stands, sample plots were selected to include the largest trees of the stand, provided the trees were not located near trails, ecotones, or other disturbances. With the largest trees possible in each 375-m<sup>2</sup> plot, the undergrowth vegetation and soil were most representative of the oldest part of the stand. Each plot was then subdivided into three 5 x 25 m macroplots. Within each 375-m<sup>2</sup> plot, all trees taller than 1 m were measured at breast height and were recorded in diameter size classes. Trees shorter than 1 m were counted in two 1 x 25 m strips along the sides of the central macroplot. In the calculation of tree basal areas, the midpoints of the diameter classes were used. Because the largest trees in the plots were included, the basal area calculation produced a higher basal area than would be expected for the average stand. The technique was used consistently throughout the study, so that the results could be compared from one stand to another.

In each stand sampled, canopy coverages were estimated for all undergrowth species within 50 2 x 5 dm microplots placed systematically along the sides of the central macroplot. The canopy coverage of each species was recorded as one of six coverage classes: 1 = <1-5%, 2 = 6-25%, 3 = 26-50%, 4 = 51-75%, 5 = 76-95%, 6 = 96-100% (Daubenmire 1959). Mosses and lichens were lumped together for coverage



estimates. Plant species in the 15 x 25 m plots which occurred in none of the 2 x 5 dm microplots were noted also.

In each stand, 25 samples of the upper decimeter of mineral soil were collected. The samples were air-dried in the field, then were composited for laboratory analysis. Soil passing through a 2-mm sieve was used to determine texture using a modified Bouyoucos method, pH using a combination electrode on the saturated soil paste, nitrogen by the Kjeldahl method, and organic matter by a modified Walkley-Black method (Moodie and Koehler 1975). Cation exchange capacity (C.E.C.), free lime, and extractable cations were determined using appropriate Hach Chemical test kits.

Nomenclature for plants collected in this study follows Van Bruggen (1976). Where possible, identification was carried out to the species level. Although plants were collected at various times during the growing season, there were some taxonomic difficulties related to the lack of flowering specimens or to the lack of clear taxonomic differences among certain species. As a result, the following were combined: (1) *Viola* spp., which includes *V. adunca*, *V. canadensis*, *V. nuttallii*, *V. pedatifida*, and *V. pratensis*; (2) Polypodiaceae, which includes *Cystopteris gracilis* and *Woodsia oregana*; (3) *Astragalus* sp., which is probably *A. adsurgens*; and (4) *Ribes setosum*, which could include *Ribes oxycanthoides*. Voucher specimens collected in this study were deposited at TRNP and in the herbarium at the University of South Dakota.

## HABITAT TYPE DESCRIPTIONS

### *Stipa comata*/*Carex filifolia* Habitat Type

This habitat type occurs on upland plateaus and gentle slopes that appear to be free of excessive erosion (fig. 3). Seven stands were sampled; 49 species were recorded (table A1). *Stipa comata*, *Carex filifolia*, *Agropyron smithii*, *Koeleria pyramidata*, and *Selaginella densa* provide the greatest coverage (table A1). *Stipa comata* forms the dominant union (Daubenmire 1968) of this association. Its abundance and stature is usually indicative of the association. During years of low moisture, *Stipa comata* produces short plants with only a few flowering culms. Close inspection of the vegetation reveals a relatively complex structure, with grasses and broadleaf forbs forming a union under *Stipa*. *Agropyron smithii* dominates this union; other conspicuous members include *Koeleria pyramidata*, *Artemisia dracunculus*, *Chrysopsis villosa*, *Lactuca oblongifolia*, *Lygodesmia juncea*, and *Tragopogon dubius*. Another union of this association is dominated by *Carex filifolia*, which has 28.5% coverage. Other important species of this low-growing union are *Artemisia frigida*, *Carex eleocharis*, *Bouteloua gracilis*, and *Gaura coccinea*. This union provides 40–50% coverage wherever it occurs in the study region. Very close to the ground surface, and in places forming dense mats, is *Selaginella densa*, which provides 47.6% coverage in this association. Outside



Figure 3.—The *Stipa comata*/*Carex filifolia* habitat type occurs on uplands where the soil is deep and well-drained. The meter stick used for scale in this and subsequent photos is painted in decimeter segments.



this association, *Selaginella densa* occurs only rarely. The *Stipa/Carex* association is the apparent climatic climax of this study region.

Major species with their constancy and mean coverage percentages are the following:

Species	Constancy	Mean coverage
<i>Stipa comata</i>	100	39.4
<i>Carex filifolia</i>	100	28.5
<i>Bouteloua gracilis</i>	86	0.9
<i>Carex eleocharis</i>	100	1.3
<i>Agropyron smithii</i>	75	10.3
<i>Koeleria pyramidata</i>	100	9.7
<i>Artemisia frigida</i>	100	3.1
<i>Gaura coccinea</i>	86	0.7
<i>Lactuca oblongifolia</i>	86	4.0
<i>Lygodesmia juncea</i>	71	0.9
<i>Tragopogon dubius</i>	100	0.4
<i>Selaginella densa</i>	100	47.6
<i>Artemisia dracunculus</i>	100	2.7

Compared to some of the other habitat types, forbs are relatively more important, although graminoids provide the greatest coverage. Among the seven stands sampled, the mean coverages of graminoids, forbs, and shrubs are the following:

graminoids: mean coverage = 90%; S.E. = 9.46  
 forbs: mean coverage = 62.4%; S.E. = 7.82  
 shrubs: mean coverage = 3.29%; S.E. = 0.71.

Forb coverage is almost 70% that of the graminoids. Hanson and Whitman (1938) reported a community in western North Dakota dominated by *Bouteloua gracilis*, *Stipa comata*, and *Carex filifolia*. They suggested this community closely reflected the macroclimate of the region. Coupland (1950, 1961) described a *Stipa comata-Bouteloua gracilis* community in western Canada in which *Carex filifolia* is important and in which *Stipa comata* is a productive species. He reported *Carex filifolia* increases in abundance in a southerly direction from Canada, and with overgrazing, *Bouteloua gracilis* becomes a dominant species. Nelson (1961) reported that heavy grazing favors the "short" grasses over the "midgrasses." In the present study, *Bouteloua gracilis* contributed little to the total canopy coverage. Coupland (1950, 1961) also found *Selaginella densa* to be an important component of the *Stipa comata*-dominated community, and suggested that *Selaginella densa* increases with protection from grazing and decreases from trampling by grazing animals on some Canadian grasslands. Mueggler and Stewart (1980) reported a *Stipa comata/Bouteloua gracilis* habitat type on alluvial substrates in southern Montana, mainly east of the Continental Divide. Their data suggest it is the most xeric steppe habitat type in western Montana. This habitat type has some similarities to the *Stipa/Carex* habitat type of TRNP, although the latter is considerably richer floristically.

Soils of this habitat type range from sandy loams to loamy sands. The high sand content of soils of *Stipa*

*comata*-dominated vegetation corresponds to other findings (Coupland 1961, Daubenmire 1970, Dix 1960, Hanson 1935, Hanson and Whitman 1938, Lauenroth and Whitman 1977, and Wright and Wright 1948). Sandy soils in this climate have a high infiltration rate, deep penetration of water, and high moisture availability, but low moisture-holding capacity. The organic matter contents of these soils were lowest of all the soils in this study and may be related to coarse textures. They readily leach, have high oxidation rates, and possibly low productivity. Edaphic characteristics are given in Appendix 3.

#### *Agropyron smithii/Carex filifolia* Habitat Type

This habitat type occurs on slopes of 9% to 20%, and is characterized by the dominance of *Agropyron smithii* and *Carex filifolia* (fig. 4). In the nine stands sampled, only 32 species occurred, compared to 49 species of the *Stipa comata/Carex filifolia* habitat type (table A2). Constancy and mean coverage percentages of the major plants of the nine stands sampled are as follows:

Species	Constancy	Mean coverage
<i>Agropyron smithii</i>	100	91.2
<i>Carex filifolia</i>	78	14.0
<i>Artemisia frigida</i>	78	1.1
<i>Tragopogon dubius</i>	78	0.2
<i>Stipa comata</i>	67	1.2
<i>Lactuca oblongifolia</i>	56	0.4

Forbs are particularly underrepresented in this habitat type. The importance of graminoids in this habitat type may be illustrated by examining the total coverage provided compared to that of the forbs and shrubs. The mean total graminoid coverage of the nine stands sampled was 111.7% (S.E. = 4.01), and the mean total forb coverage in the same nine stands was 2.33% (S.E. = 0.91). Shrubs also provided little coverage in these stands; mean total shrub coverage was 1.44% (S.E. = 0.38).

The soils of this habitat type are fine-textured. Hanson and Whitman (1938) described a plant community in western North Dakota dominated by *Agropyron smithii*, *Bouteloua gracilis*, and *Carex filifolia*, which occurred on gradual slopes of heavy-textured soils. Quinnild and Cosby (1958) found vegetation of mesa tops in western North Dakota dominated by *Agropyron smithii*, *Agropyron dasystachyum*, *Stipa comata*, *Bouteloua gracilis* and *Artemisia frigida*. Larson and Whitman (1942) also reported on *Agropyron smithii*-dominated vegetation on mesas in the badlands of South Dakota. Their results suggest protection maintains a vegetation type dominated by *Agropyron smithii* and *Carex* sp., while combined mowing and grazing produces vegetation dominated by *Bouteloua gracilis*.

All surface soils of this habitat type were classified as loams; the clay content ranged from 14.0% to 24.4%. Fine-textured soils in this semiarid region have poor water balance conditions. This may account for the less





Figure 4.—Long slope of fine-textured soil which supports the *Agropyron smithii*/*Carex filifolia* habitat type.

rich assortment of species though *Agropyron smithii* is favored and often occurs in dense stands on heavy soils in this region (Weaver and Albertson 1956). The *Agropyron*/*Carex* association at TRNP is an edaphic climax.

#### ***Andropogon scoparius*/*Carex filifolia* Habitat Type**

Nine stands of this habitat type were sampled. A total of 66 species, including nine shrub species, were found. This habitat type is characterized by the dominance of *Andropogon scoparius* (fig. 5) with a mean coverage of 76.8% (table A3).

Characteristic species of this habitat type with constancy and mean coverage percentages are the following:

Species	Constancy	Mean coverage
<i>Andropogon scoparius</i>	100	78.8
<i>Calamovilfa longifolia</i>	100	5.3
<i>Carex filifolia</i>	100	8.8
<i>Koeleria pyramidata</i>	89	0.7
<i>Echinacea angustifolia</i>	89	0.6
<i>Aster oblongifolius</i>	89	0.4
<i>Artemisia frigida</i>	89	0.4
<i>Helianthus rigidus</i>	78	3.6
<i>Lactuca oblongifolia</i>	67	1.0
<i>Gaura coccinea</i>	67	0.5
<i>Lygodesmia juncea</i>	67	0.2

Although forbs provide limited coverage, there are numerous species which add considerable diversity to

the vegetation of this habitat type. The mean coverages of graminoids, forbs, and shrubs in stands of this habitat type are as follows:

graminoids:	mean coverage = 97.4%;	S.E. = 2.86
forbs:	mean coverage = 7.78%;	S.E. = 2.59
shrubs:	mean coverage = 1.22%;	S.E. = 0.41.

Although more forb and shrub species are present in this habitat type, both life forms provide much less coverage than in the *Stipa*/*Carex* habitat type.

Because of the topographic position and the coarse-textured soils generally associated with the stands studied, the vegetation is considered to be a topoedaphic climax.

Hanson and Whitman (1938) described an *Andropogon scoparius*-dominated community on northerly slopes and where snow accumulates. They observed that *Andropogon* colonized bare areas created by step erosion. Once established, it spread and continued to dominate larger eroded areas, and its density and height made it an effective competitor with shorter species. The stands of *Andropogon*/*Carex* habitat type studied also may have become established after erosion. They appear to be permanent; there is no evidence other species are replacing the *Andropogon*. Redmann (1975) described a community dominated by *Andropogon scoparius*. His stands were on north-facing slopes, and both *Helianthus rigidus* and *Rosa arkansana* were important species.





Figure 5.—Well-developed stand of *Andropogon scoparius*/*Carex filifolia* association.

Both Hanson and Whitman (1938) and Redmann (1975) observed small stands of *Andropogon gerardii*-dominated vegetation. The present study also found small stands of vegetation, usually on steep south-facing slopes and primarily in the North Unit. These stands are small and are not considered sufficiently large to sample or to classify as a habitat type.

Soils of this habitat type have high sand and low clay contents. In three of the stands studied, soils had considerable gravel, from 7% to 36% by weight. In the South Unit of TRNP, the five stands sampled were on soils rich in gravel or scoria, on slopes ranging from 18% to 42%. In the North Unit of TRNP, this habitat type is restricted primarily to the glaciated areas north of the Missouri River, though a few stands were observed south of the River, in the Achenbach Hills region. Stands 22, 23, and 24 had gravel ranging from 6.7% to 36.2% of the soil dry weight. Edaphic characteristics are given in Appendix 3.

#### ***Juniperus horizontalis*/*Andropogon scoparius* Habitat Type**

Five stands of this vegetation were sampled; all are in the South Unit of the Park and on slopes exceeding 28%. This habitat type is distinguished by the dominance of *Juniperus horizontalis*, with a mean coverage of 80.4%, and the presence of the *Andropogon scoparius* union (fig. 6). The *Carex filifolia* union is also represented (table A4). In the stands sampled, there were 60 species, 43 of which also occur in the *A. scoparius*/*C. filifolia*

habitat type (table A4). Based on a floristic comparison, the *Andropogon*/*Carex* and *Juniperus*/*Andropogon* habitat types are 68% similar. Ten shrub species occur in the *Juniperus*/*Andropogon* habitat type, although the coverage of *J. horizontalis* exceeds considerably the combined coverage of all the other shrubs. Stand 10 was richest with eight shrub species, including *J. horizontalis*, *Artemisia cana*, *Potentilla fruticosa*, *Prunus virginiana*, *Rhus aromatica*, *Rosa arkansana*, *Shepherdia argentea*, and *Symphoricarpos occidentalis*. There, the coverage of *J. horizontalis* was 83%, and the total coverage of the remaining seven shrubs was 30%. This habitat type is rather localized in its distribution. On the basis of topographic and edaphic factors, vegetation of this habitat type is a topoedaphic climax. Major species with their constancy and mean coverage percentages in this association are as follows:

Species	Constancy	Mean coverage
<i>Juniperus horizontalis</i>	100	80.4
<i>Andropogon scoparius</i>	100	24.2
<i>Calamovilfa longifolia</i>	100	7.0
<i>Carex filifolia</i>	100	3.0
<i>Campanula rotundifolia</i>	100	1.1
<i>Petalostemon purpureum</i>	100	0.6
<i>Galium boreale</i>	80	4.8
<i>Anemone patens</i>	80	2.4
<i>Potentilla fruticosa</i>	80	2.4
<i>Gaura coccinea</i>	80	0.6
<i>Echinacea angustifolia</i>	80	0.5
<i>Linum perenne</i>	80	0.3
<i>Senecio plattensis</i>	80	0.2





Figure 6.—*Juniperus horizontalis*/*Andropogon scoparius* habitat type. The low-growing *Juniperus* is evident as a low growing, darker plant in this photo. Scattered culms of *Andropogon* are also conspicuous.

Of these 13 species with high constancies, only five have 67% or higher constancies in the *Andropogon*/*Carex* association. Comparing the lists of species with high constancy values for the *Andropogon*/*Carex* and *Juniperus*/*Andropogon* associations, the floristic similarity is only 38%, considerably lower than the similarity based on the total species lists of both associations. Coupland (1961) reported *J. horizontalis* on exposed sites in the Great Sand Hills, Alberta. Quinnild and Cosby (1958) found it on eroded edges of a mesa in western North Dakota. Whitman and Hanson (1939) described a community on scoria buttes in North Dakota, in which both *J. horizontalis* and *Mentzelia decapetala* were dominant. Redmann (1975) observed occasional mats of *J. horizontalis* on sandy soil in western North Dakota. At TRNP, stands of *J. horizontalis* appear to be stable, and habitat type status is assigned to these sites. Edaphic characteristics of this habitat type are listed in Appendix 3.

#### ***Artemisia tridentata*/*Agropyron smithii* Habitat Type**

This habitat type occurs on terraces or narrow benches that parallel the contours of hills 10 to 60 m above the valley floor of the Little Missouri River and its principal tributaries (fig. 7). In the four stands of this

vegetation, only 29 species were found. *Artemisia tridentata* is dominant, and both the *Agropyron smithii* and *Carex filifolia* unions are represented (table A5). There are nine shrub species, but the coverage of *A. tridentata* far exceeds the combined coverage of the remaining eight shrubs. *Atriplex confertifolia* is second, with slightly less than 2% mean coverage. The paucity of species reflects the severity of the habitats supporting this vegetation, which is also a topoedaphic climax. Major species of this association and their constancy and mean coverage percentages are as follows:

Species	Constancy	Mean coverage
<i>Artemisia tridentata</i>	100	29.7
<i>Agropyron smithii</i>	100	68.8
<i>Agropyron caninum</i>	100	8.1
<i>Opuntia polyacantha</i>	100	0.4
<i>Atriplex confertifolia</i>	75	1.5
<i>Carex filifolia</i>	50	3.2

All species of the four stands studied are listed in table A5. Shrubs and graminoids are most conspicuous in this habitat type; forbs are sparse. The mean coverages of shrubs, graminoids, and forbs are as follows:

shrubs:	mean coverage = 33.5%;	S.E. = 4.35
graminoids:	mean coverage = 83.0%;	S.E. = 8.83
forbs:	mean coverage = 3.5%;	S.E. = 1.71.





Figure 7.—*Artemisia tridentata*/*Agropyron smithii* habitat type This habitat type occurs on narrow benches or terraces along the contours of the hills.

Edaphic characteristics of the four stands studied are listed in Appendix 3.

Other descriptions of *Artemisia tridentata*-dominated vegetation of western North Dakota and eastern Montana indicate broad similarities in composition and habitat conditions (Brown 1971, Hazlett and Hoffman 1975, Whitman and Hanson 1939). In the Missouri River "breaks" region of Montana, Mackie (1970) described an *Artemisia tridentata*/*Agropyron smithii* association as an edaphic climax on "shallow, gravelly, or claypan surface soils."

In his study of badland vegetation in southeastern Montana, Brown (1971) described a community dominated by *Artemisia tridentata*, with *Atriplex confertifolia* and *Agropyron spicatum* as codominants. *Agropyron smithii* was present in small patches.

#### ***Artemisia cana*/*Agropyron smithii* Habitat Type**

This habitat type occurs on large areas of floodplains and river terraces adjacent the Little Missouri River and its tributaries, areas commonly referred to as "sagebrush flats" (fig. 8). The dominant is *Artemisia cana*, with a mean coverage of 33.6% (table A6). Between widely spaced shrubs, numerous other species occur in one or two lower unions dominated by *Agropyron smithii* and *Carex filifolia*. Although 55 species occurred in the 12 stands sampled, only seven have a constancy of 50% or higher. Major species of this

association and their constancy and mean coverage percentages are as follows:

Species	Constancy	Mean coverage
<i>Artemisia cana</i>	100	33.6
<i>Agropyron smithii</i>	100	70.9
<i>Stipa viridula</i>	75	2.4
<i>Agropyron caninum</i>	67	6.6
<i>Artemisia frigida</i>	58	0.6

As in the *Artemisia tridentata*/*Agropyron smithii* habitat type, graminoids in this habitat type provide considerably more coverage than either forbs or shrubs. The total forb and shrub coverage in this habitat type, however, is greater than in the *A. tridentata*/*A. smithii* habitat type. The means for this habitat type are the following:

shrubs:	mean coverage = 54.1%;	S.E. = 7.69
graminoids:	mean coverage = 84.8%;	S.E. = 5.32
forbs:	mean coverage = 9.4%;	S.E. = 2.27.

Floristically, this habitat type is only 47% similar to that of the *A. tridentata*/*A. smithii* habitat type. *Symphoricarpos occidentalis* occurs in five of the stands sampled; its mean coverage is 47%. Clones of this species are described in detail later. *Agropyron caninum* and *Muhlenbergia cuspidata* reach their maximum abundance in this and the *A. tridentata*/*A. smithii* habitat type.





Figure 8a.—*Artemisia cana*/*Agropyron smithii* habitat type. The relatively wide spacing of *Artemisia* and nearly complete herbaceous cover is evident in this photo.



Figure 8b.—Broad level deposit of alluvium support *Artemisia cana*-dominated vegetation. These areas are referred to as "sagebrush flats."



This vegetation is heavily utilized by bison and deer in TRNP. Nelson (1961) suggested that heavy grazing of this vegetation reduces the amounts of *A. smithii* and *Stipa viridula* in favor of *Bouteloua gracilis*. Hanson and Whitman (1938) suggested that occasional flooding and deposition of alluvium might prevent short grasses from becoming important in this vegetation. *A. cana* provides fall and winter browse for game animals and, outside TRNP, for livestock (Johnson and Nichols 1970).

The substrates vary considerably with respect to drainage and texture, and have moderate salt content (Hanson and Whitman 1938). The *A. cana/A. smithii* association is an edaphic climax in this region.

This is an important habitat type, covering large areas. Outside the Park, grazing animals utilize it heavily, and ungrazed stands of the habitat type inside the Park become even more valuable as reference areas. Edaphic characteristics are listed in Appendix 3.

#### ***Fraxinus pennsylvanica/Symphoricarpos occidentalis* Habitat Type**

In the arid northern Great Plains, woodland vegetation is confined to habitats where adequate soil moisture, along streams or in other favorable topographic positions, compensates for the arid climate (fig. 9). The

*F. pennsylvanica/S. occidentalis* association is an edaphic climax on the floodplain adjacent the Little Missouri River and its major tributaries. Tree population data for this habitat type are given in Appendix 2. *Populus deltoides* currently dominates many of the stands but is no longer reproducing. The data indicate it will be replaced by *Fraxinus pennsylvanica*. Tree basal areas of the stands studied range from 23.3 m<sup>2</sup>/ha to 63.1 m<sup>2</sup>/ha (Appendix 2). The larger trees, some 6–7 dm d.b.h., are *Populus deltoides*, but the young trees establishing in the stands are *Fraxinus pennsylvanica*. *Juniperus scopulorum* is tallied with the tree species in the data, although it is an understory species in the closed forest. Its current abundance is attributed to adequate light penetrating to the shrub and herb layers of the community as a result of wide spacing of the old *Populus*. Along the Missouri River, in central North Dakota (Burgess et al. 1973), in southeastern South Dakota (Wilson 1970), and near Omaha, Nebr. (Vaubel 1975), *Populus deltoides* is a pioneer species, and is replaced successional by various combinations of *Fraxinus*, *Ulmus*, *Acer*, and *Celtis*. The *F. pennsylvanica/S. occidentalis* habitat type is recognized by the combination of tree and undergrowth unions. *F. pennsylvanica* is present in the understory, although the overstory may be dominated currently by the non-reproducing *Populus deltoides*. The undergrowth is dominated by the Sym-



Figure 9.—The *Fraxinus pennsylvanica/Symphoricarpos occidentalis* habitat type is confined in this semiarid climate to relatively narrow bands paralleling larger streams. Along a meander of the Little Missouri River, shown here, *Populus deltoides* became established in rows parallel to the stream course.



*phoricarpus occidentalis* union with *S. occidentalis*, *Rhus aromatica*, *Toxicodendron rydbergii*, and other shrubs well-represented (fig. 10). Among the grasses, *Calamovilfa longifolia*, *Elymus canadensis*, and *Muhlenbergia racemosa* are important (table A7). In the four stands sampled, there are 52 undergrowth species. Major undergrowth species of the stands sampled and their constancy and mean coverage percentages are as follows:

Species	Constancy	Mean coverage
<i>Symphoricarpos occidentalis</i>	100	40.5
<i>Toxicodendron rydbergii</i>	100	23.0
<i>Melilotus officinalis</i>	100	28.8
<i>Elymus canadensis</i>	100	11.5
<i>Muhlenbergia racemosa</i>	100	8.3
<i>Melilotus albus</i>	100	7.0
<i>Rhus aromatica</i>	100	5.4
<i>Agropyron caninum</i>	100	4.0
<i>Smilacina stellata</i>	100	0.9
<i>Parthenocissus vitacea</i>	100	0.6
<i>Poa pratensis</i>	75	37.2

The importance of shrubs, graminoids, and forbs may be shown by their mean coverage values:

shrubs:	mean coverage = 76.8%;	S.E. = 23.2
graminoids:	mean coverage = 64.2%;	S.E. = 27.0
forbs:	mean coverage = 43.5%;	S.E. = 15.8.

*Symphoricarpos* and *Toxicodendron* together provide 63.5% coverage among the shrubs.

Before TRNP excluded cattle, grazing was common in these streamside forests. Currently, in TRNP, bison sometimes utilize this habitat type for grazing, watering, and the shade provided during the summer. The importance of certain alien species attests to the heavy usage earlier by cattle, and the considerable usage by bison now that the area is a National Park. The most important grass is *Poa pratensis*, with a mean coverage of 36.2%. The most important forb is *Melilotus officinalis*, with a mean coverage of 28.8%. Edaphic characteristics of stands sampled are listed in Appendix 3.

#### *Fraxinus pennsylvanica*/*Prunus virginiana* Habitat Type

This habitat type occurs in ravines or draws or on moderately steep north-facing slopes throughout much of TRNP. It is recognized by the presence of *Fraxinus pennsylvanica* as the dominant species, with *Ulmus americana* as a codominant in some of the stands. Both species reproduce successfully in this habitat type. In V-shaped ravines, largest *Fraxinus* occur near the center, or bottom, of the stand, where there is greater soil moisture. Tree basal area of the stands studied ranged from 21.6 m<sup>2</sup>/ha to 35.9 m<sup>2</sup>/ha (Appendix 2). The largest tree sampled was a *Fraxinus*, with 116 xylem rings at breast height. The average age of the *Fraxinus*



Figure 10.—Interior of a stand of the *Fraxinus pennsylvanica*/*Symphoricarpos occidentalis* habitat type. The undergrowth here is a rich mixture of both shrubs and herbaceous species, with *Fraxinus* seedlings obvious in the foreground.



was 60–70 years. Heart rot prevented determining the age of some of the larger *Fraxinus*. *Ulmus americana* is important in some stands, and *Acer negundo* and *Juniperus scopulorum* also occur in this habitat type. The undergrowth is dominated by *Prunus virginiana*, 2–3 m tall, with a mean coverage of 27.7% (fig. 11). This shrub reaches its greatest density near the edge of these stands, where light intensity is greater, and it often forms a fringe beyond the stand margin. The smaller *Symphoricarpos occidentalis* and *Rosa woodsii* are also important shrubs in the undergrowth (table A7). The important graminoids of this habitat type are *Carex sprengelii*, with a mean coverage of 61.1%, and *Elymus virginicus*, with a mean coverage of 27.5%. These two species reach their maximum importance in this habitat type. Owing to heavy utilization by cattle in the past, and possibly as a result of bison use currently, *Poa pratensis* and *Taraxacum officinale* are both important in this habitat type. Tree seedlings and saplings are abundant; trampling by bison apparently is not an important factor at present. *Galium boreale*, with a mean coverage of 10.2%, is the most important forb. Other forbs characteristic of this habitat type include *Monarda fistulosa*, *Thalictrum dasycarpum*, *T. venulosum*, *Sanicula marilandica*, *Arctium minus*, and *Ranunculus arbor-tivus*. In the four stands sampled, 51 species were found. The vegetation appears to be a topographic climax. Important undergrowth species in stands of this habitat

type and their constancy and mean coverage percentages are as follows:

Species	Constancy	Mean coverage
<i>Prunus virginiana</i>	100	27.7
<i>Symphoricarpos occidentalis</i>	100	12.1
<i>Elymus virginicus</i>	100	27.5
<i>Galium boreale</i>	100	10.2
<i>Carex sprengelii</i>	50	30.6
<i>Poa pratensis</i>	75	15.6

Mean coverages for shrubs, graminoids, and forbs are as follows:

shrubs:	mean coverage = 46.2%;	S.E. = 14.0
graminoids:	mean coverage = 82.2%;	S.E. = 19.0
forbs:	mean coverage = 29.0%;	S.E. = 11.6.

Compared to the *Fraxinus*/*Symphoricarpos* habitat type, graminoids provide greater coverage, but both shrubs and forbs provide less coverage. Overall, the total mean undergrowth coverage is 157.4% for the *Fraxinus*/*Prunus* habitat type, compared to 184.5% for the *Fraxinus*/*Symphoricarpos* habitat type. Stands of this habitat type are important deer habitats, providing both browse and shelter. Stands also attract domestic grazing animals outside the boundary of TRNP. Lack of tree



Figure 11.—*Fraxinus pennsylvanica*/*Prunus virginiana* habitat type. Photo taken from the edge of the stand. *P. virginiana* dominates the undergrowth.



reproduction in such stands have concerned other investigators (Boldt et al. 1978, Severson and Boldt 1978). It appears from their studies, and also from results of a study in the Custer National Forest, that cattle grazing and trampling is the most important single cause of failure of trees to successfully replace themselves in these wooded draws. The stands in TRNP become all the more important, therefore, as reference areas with which to compare changes resulting from domestic animal activity outside the Park. Even within the Park, considerable variation in undergrowth vegetation among stands of this habitat type reflect the effects of grazing before the area was a national park.

Edaphic characteristics of stands of this habitat type are given in Appendix 3.

#### ***Populus tremuloides*/Betula occidentalis Habitat Type**

This habitat type occurs on upper slopes facing northwest to east (fig. 12). Stands of the *Fraxinus*/*Prunus* habitat type are lower on the same slopes. The habitat type is not widespread in TRNP, and is recognized by the dominance of *Populus tremuloides*. The vegetation is also a topographic climax. The undergrowth is dominated by *Betula occidentalis*, with a mean coverage of 22.1%, *Prunus virginiana*, with a mean coverage of 31.1%, and *Symphoricarpos albus*, with a mean coverage of 20.5%. *S. occidentalis* and *Toxicodendron*

*rydbergii* are also important undergrowth shrubs in this habitat type. *Carex sprengelii*, *Oryzopsis micrantha*, and *Apocynum androsaemifolium* are major herbaceous species in this habitat type, with *Disporum trachycarpum*, *Galium boreale*, and *Smilacina stellata* also important. In the four stands sampled, 57 species were encountered (table A7). Mean basal area of trees in stands sampled is 24.6 m<sup>2</sup>/ha. The largest size class represented among the tree species is 2–3 dm d.b.h. The largest *Populus*, based on xylem layers at breast height, ranged from 43 to 77 years. Major undergrowth species in stands of this habitat type and their constancy and mean coverage percentages are as follows:

Species	Constancy	Mean coverage
<i>Betula occidentalis</i>	100	22.1
<i>Prunus virginiana</i>	100	31.1
<i>Toxicodendron rydbergii</i>	100	22.7
<i>Disporum trachycarpum</i>	100	5.4
<i>Smilacina stellata</i>	100	3.6
<i>Galium boreale</i>	100	5.0
<i>Apocynum androsaemifolium</i>	75	11.4

Graminoids provided less coverage in this habitat type than in any other habitat type studied. Mean coverages for shrubs, graminoids, and forbs in stands sampled are as follows:



Figure 12.—*Populus tremuloides*/Betula occidentalis habitat type. Undergrowth is a rich mixture of shrubs and forbs primarily.



shrubs: mean coverage = 91.5%; S.E. = 14.2  
 graminoids: mean coverage = 20.8%; S.E. = 9.1  
 forbs: mean coverage = 43.2%; S.E. = 8.9.

Shrub coverage is higher than that of other habitat types, and forb coverage is the same as that of the *Fraxinus/Symphoricarpos* habitat type but higher than that of the *Fraxinus/Prunus* habitat type. There is a 46% floristic similarity between stands of this habitat type and those of the *Fraxinus/Prunus* habitat type. The *Fraxinus/Prunus* and *Populus/Betula* habitat types are in relatively close proximity; and although they share nearly half their species, the two habitat types are distinct and can be separated on the basis of critical overstory and undergrowth species.

As shown in Appendix 2, the tree stratum in stands of this habitat type has six species, although seldom do all occur in the same stand. Both *P. tremuloides* and *F. pennsylvanica* have population structures indicating successfully reproducing species that could maintain or increase their population sizes. *Fraxinus* is currently a codominant in this habitat type; but it may become a more important species in the future. A constant source of seeds is available from lower slope stands of the *Fraxinus/Prunus* habitat type. Edaphic characteristics are given in Appendix 3.

#### ***Juniperus scopulorum/Oryzopsis micrantha* Habitat Type**

This habitat type occurs on northwest to north-facing slopes of 35–70% and is recognized by the dominance of *J. scopulorum*; the vegetation is a topographic climax. Stands of this habitat type ordinarily cover entire hill-sides in TRNP, making them quite distinct even from a distance (fig. 13). Principal shrubs are *Prunus virginiana*, *Symphoricarpos occidentalis*, and *Rhus aromatica*; their abundance relates positively with openings among the dominant *Juniperus*. *Oryzopsis micrantha* dominates the undergrowth and is also characteristic of this habitat type. Mosses and lichens cover about 72% of the ground surface (table A8). In the seven stands of this habitat type sampled, 52 species occurred.

Major undergrowth species in stands of this habitat type and their constancy and mean coverage percentages are as follows:

Species	Constancy	Mean coverage
<i>Oryzopsis micrantha</i>	100	66.9
Mosses and Lichens	100	72.3
<i>Galium boreale</i>	100	4.5
<i>Rhus aromatica</i>	100	4.2
<i>Symphoricarpos occidentalis</i>	100	1.7
<i>Campanula rotundifolia</i>	100	1.0
<i>Prunus virginiana</i>	86	10.5

Mean coverages provided by all shrubs, graminoids, and forbs are the following:

shrubs: mean coverage = 17.4%; S.E. = 7.76  
 graminoids: mean coverage = 69.1%; S.E. = 3.72  
 forbs: mean coverage = 9.4%; S.E. = 1.77.

The mean forb coverage is calculated without the moss and lichen coverage. Shrubs are not as important in this association as in the other forest associations of TRNP. The undergrowth is dominated by herbaceous species primarily.

*Fraxinus pennsylvanica* is present in most of the stands; but most are seedlings which grow in slight depressions where soil moisture exceeds normal for this habitat type. There is no evidence at present to suggest that *Fraxinus* will replace *Juniperus scopulorum* as the climax dominant. The largest *Juniperus* sampled are in the 3–4-dm d.b.h. size class (appendix 2). Occasionally, larger ones are observed. Many of the trees have more than 125 xylem layers, and some have more than 250. Tree basal areas for the stands sampled ranged from 22.1 m<sup>2</sup>/ha to 29.6 m<sup>2</sup>/ha. Before TRNP was established, the primary use made of these stands was the harvest of *Juniperus* for use as fence posts (Ralston 1960). Limited cattle grazing occurred, although game species used the stands extensively. Deer still utilize stands of this habitat type quite extensively. There are numerous game trails over much of the area of this habitat type; *Achillea millefolium* is most abundant along these trails. The physiognomy of these stands is similar to that of the pinyon-juniper stands of the Southwest, although pine is absent. Edaphic characteristics are given in Appendix 3.

#### ***Symphoricarpos occidentalis* Community**

This community occurs as scattered thickets over much of the study area. It is recognized by the dominance of *Symphoricarpos occidentalis* which forms dense thickets a few to several meters across (fig. 14). Observations suggest that these thickets may be long-lived seral communities (associates) occurring in at least two habitat types in TRNP. Typically, the community has three layers, an upper layer 0.5–1.0 m tall dominated by *Symphoricarpos occidentalis* and *Rosa woodsii*, an intermediate layer 0.2–1.5 m tall dominated by *Agropyron smithii*, *A. caninum*, *Lactuca oblongifolia*, and *Poa pratensis*, and a lower layer 0.1–0.3 m tall dominated by *Achillea millefolium*, *Artemisia ludoviciana*, and *Galium boreale*. In some of the larger, more complex thickets, there is a fourth taller layer dominated by *Prunus virginiana*, 2–3 m tall, with *Ribes odoratum* and *Rosa woodsii*. A total of 37 species occur in the five stands sampled. Most of the forbs grow near the margins of the thickets, where the shrubs are less dense. *Artemisia ludoviciana* and *Parietaria pennsylvanica* reach their maximum coverage and constancy in this community.

*Symphoricarpos occidentalis* thickets at TRNP occur on various exposures and substrates, from fine-textured alluvial soils of the *Artemisia cana/Agropyron smithii* habitat type to the coarse, gravelly substrates of the *Andropogon scoparius/Carex filifolia* habitat type. They also occur near margins of *Fraxinus*-dominated habitat types described above.

*Symphoricarpos* thickets also have been described in steppe vegetation elsewhere (Daubenmire 1970, Mackie 1970).





Figure 13a—Stand of *Juniperus scopulorum*/*Oryzopsis micrantha* habitat type on north-facing slope.



Figure 13b.—Interior of stand of *Juniperus*/*Oryzopsis* association showing the characteristically dense vegetation.





Figure 14.—Dense thicket of *Symphoricarpos occidentalis* with others in the background.

### KEY TO THE HABITAT TYPES

1. Tree species absent
2. Herbaceous species dominant
  3. *Stipa Comata* well represented; *Agropyron smithii* and/or *Andropogon scoparius* inconspicuous or absent ..... *Stipa comata*/*Carex filifolia* habitat type.
  3. *Agropyron smithii* and/or *Andropogon scoparius* dominant; *Stipa comata* inconspicuous or absent.
    4. *Agropyron smithii* abundant; *Andropogon scoparius* absent ..... *Agropyron smithii*/*Carex filifolia* habitat type.
    4. *Andropogon scoparius* abundant ..... *Andropogon scoparius*/*Carex filifolia* habitat type.
2. Shrub species dominant
  5. *Artemisia tridentata* or *A. cana* present and conspicuous.
    6. *Artemisia tridentata* dominant; *Artemisia cana* absent ..... *Artemisia tridentata*/*Agropyron smithii* habitat type.
    6. *Artemisia cana* dominant; *Artemisia tridentata* absent ..... *Artemisia cana*/*Agropyron smithii* habitat type.
  5. *Artemisia tridentata* and *A. cana* absent or inconspicuous.
    7. *Juniperus horizontalis* principal shrub; *Symphoricarpos occidentalis* if present is widely scattered and dwarfed ..... *Juniperus horizontalis*/*Andropogon scoparius* habitat type.
    7. *Symphoricarpos occidentalis* forming dense thickets or clones up to 1 m or more in height ..... *Symphoricarpos occidentalis* community type.
1. Tree species present and dominant
  8. *Populus deltoides* and/or *Fraxinus pennsylvanica* dominant or exhibiting adequate reproduction to become dominant.
    9. Undergrowth dominated by *Prunus virginiana*... ..... *Fraxinus pennsylvanica*/*Prunus virginiana* habitat type.
    9. Undergrowth dominated by *Symphoricarpos occidentalis*; *Prunus virginiana* absent or inconspicuous ..... *Fraxinus pennsylvanica*/*Symphoricarpos occidentalis* habitat type.
  8. *Populus deltoides* and/or *Fraxinus pennsylvanica* absent or present in seedling and sapling size only.
    10. *Populus tremuloides* dominant; *Juniperus scopulorum* absent or rare..... ..... *Populus tremuloides*/*Betula occidentalis* habitat type.
    10. *Juniperus scopulorum* dominant; *Populus tremuloides* absent ..... *Juniperus scopulorum*/*Oryzopsis micrantha* habitat type.

## DISCUSSION

### Validation of the Habitat Type Classification

Natural vegetation is a convenient characteristic of the landscape to use in a classification scheme; the more mature the vegetation is, the more closely it reflects the biotic potential of the landscape (Daubenmire 1976).

A classification scheme provides a systematic ordering of the landscape units under study. In this case, the units are the habitat types, and the manner of delimiting these provides as natural a classification as possible. Because a habitat type consists of land units representing areas of similar environment, other research efforts can be related to the habitat types. For example, wildlife utilization (Mackie 1970), small mammal distributions (Hoffman 1960, Rickard 1957), growth rates of ponderosa pine (Daubenmire 1961), depth of soil moisture drying (Daubenmire 1972), and productivity and grazing potentials (Bjugstad and Whitman 1970, Mueggler and Stewart 1980), can be related to the habitat types to provide a means of correlating and communicating research results. Silvicultural and watershed management implications also have been discussed in relation to habitat types (Hoffman and Alexander 1976, 1980, 1983).

### Species Diversity

Species diversity may be described as the median number of species occurring in areas of constant sam-

pling size (Daubenmire 1970). Species diversity of the habitat types encountered in this study is shown in table 2. Among the steppe habitat types, diversity ranged from a low of 10 in the *Agropyron smithii*/*Carex filifolia* habitat type to a high of 29 in the *Juniperus horizontalis*/*Andropogon scoparius* habitat type. Four of the five stands sampled of the *Juniperus*/*Andropogon* habitat type are on north-facing slopes, with reduced solar insolation. This may be an important factor in influencing species diversity in the semi-arid climate of western North Dakota. Among the edaphic characteristics measured, surface soils of this habitat type are similar to those of the *Andropogon scoparius*/*Carex filifolia* and *Agropyron smithii*/*Carex filifolia* habitat types, the latter of which had the lowest median species diversity. The low species diversity of the *Agropyron*/*Carex* habitat type is possibly related to the very high coverage of *Agropyron* (table A2). Shantz (1924) described dense, almost pure stands of *Agropyron smithii*, accompanied by a paucity of forbs in west-central South Dakota. Possibly the density of the *Agropyron* sod inhibits the successful establishment of numerous other species.

Species diversity among habitat types dominated by tree species ranged from 21 in the *Juniperus scopulorum*/*Oryzopsis micrantha* habitat type to 30 in the *Populus tremuloides*/*Betula occidentalis* habitat type, which included six tree species. The remaining two habitat types dominated by trees had 29 species each. The generally higher numbers of species in the tree-dominated habitat types probably reflects the wetter conditions compared to the steppe vegetation of TRNP.

Table 2.—Species diversity in habitat types of Theodore Roosevelt National Park

Habitat type	No. of stands studied	Median no. of shrub, graminoid, and forb species per H.T. <sup>1</sup>	Total tree species per H.T. <sup>2</sup>
<i>Stipa comata</i> / <i>Carex filifolia</i>	7	23	
<i>Agropyron smithii</i> / <i>Carex filifolia</i>	9	10	
<i>Andropogon scoparius</i> / <i>Carex filifolia</i>	9	22	
<i>Juniperus horizontalis</i> / <i>Andropogon scoparius</i>	5	29	
<i>Artemisia tridentata</i> / <i>Agropyron smithii</i>	4	14	
<i>Artemisia cana</i> / <i>Agropyron smithii</i>	12	13	
<i>Fraxinus pennsylvanica</i> / <i>Symphoricarpos occidentalis</i>	4	26	3
<i>Fraxinus pennsylvanica</i> / <i>Prunus virginiana</i>	4	25	4
<i>Populus tremuloides</i> / <i>Betula occidentalis</i>	4	24	6
<i>Juniperus scopulorum</i> / <i>Oryzopsis micrantha</i>	7	19	2
<i>Symphoricarpos occidentalis</i> community type	5	15	

<sup>1</sup>Based on fifty 0.1 m<sup>2</sup> microplots, per stand.

<sup>2</sup>Based on 375 m<sup>2</sup> per stand.



## LITERATURE CITED

- Arno, Stephen F., and Robert D. Pfister. 1977. Habitat types: An improved system for classifying Montana's forests. *Western Wildlands* (Spring) 3(4):6-11.
- Bjurgstad, Ardell J., and Warren C. Whitman. 1970. Significance of reduced plant vigor in relation to range condition. *Journal of Range Management* 23(3):181-184.
- Boldt, Charles D., Daniel W. Uresk, and Kieth E. Severson. 1978. Riparian woodlands in jeopardy on Northern High Plains. p. 184-189. In *Strategies for protection and management of floodplain wetlands and other riparian ecosystems*. USDA Forest Service General Technical Report WO-12, 410 p. Washington, D.C.
- Brown, Ray W. 1971. Distribution of plant communities in southeastern Montana badlands. *American Midland Naturalist* 85:458-477.
- Burgess, Robert L., W. Carter Johnson, and Warren R. Keammerer. 1973. Vegetation of the Missouri River floodplain in North Dakota. Report to the Office of Water Resources Research, 162 p. U.S. Department of Interior, Washington, D.C.
- Coupland, Robert T. 1950. Ecology of mixed prairie in Canada. *Ecological Monographs* 20:271-315.
- Coupland, Robert T. 1961. A reconsideration of grassland classification in the northern Great Plains of North America. *Journal of Ecology* 49:135-167.
- Daubenmire, R. 1959. A canopy-coverage method of vegetation analysis. *Northwest Science* 33:43-66.
- Daubenmire, R. 1961. Vegetative indicators of rate of height growth in ponderosa pine. *Forest Science* 7:24-34.
- Daubenmire, R. 1968. Plant communities. A textbook of plant synecology. 300 p. Harper and Row, New York, N.Y.
- Daubenmire, R. 1970. Steppe vegetation of Washington. Technical Bulletin 62, 131 p. Washington Agriculture Experiment Station, Washington State University, Pullman.
- Daubenmire, R. 1972. Annual cycles of soil moisture and temperature as related to grass development in the steppe of eastern Washington. *Ecology* 53:419-424.
- Daubenmire, R. 1976. The use of vegetation in assessing the productivity of forest lands. *Botanical Review* 42:115-143.
- Daubenmire, R., and Jean B. Daubenmire. 1968. Forest vegetation of eastern Washington and northern Idaho. Technical Bulletin 60, 104 p. Washington Agriculture Experiment Station, Washington State University, Pullman.
- Dix, Ralph L. 1960. The effects of burning on the mulch structure and species composition of grasslands in western North Dakota. *Ecology* 41:49-56.
- Hanson, Herbert C. 1935. A comparison of methods of botanical analysis of the native prairie in western North Dakota. *Journal of Agricultural Research* 49:815-842.
- Hanson, Herbert C., and Warren C. Whitman. 1938. Characteristics of major grassland types in western North Dakota. *Ecological Monographs* 8:57-114.
- Hazlett, Donald L., and George R. Hoffman. 1975. Plant species distribution patterns in *Artemisia tridentata* and *Artemisia cana*-dominated vegetation in western North Dakota. *Botanical Gazette* 136:72-77.
- Hoffman, George R. 1960. The small mammal components of six climax plant associations in eastern Washington and northern Idaho. *Ecology* 41:571-575.
- Hoffman, George R., and Robert R. Alexander. 1976. Forest vegetation of the Bighorn Mountains, Wyoming: A habitat type classification. USDA Forest Service Research Paper RM-170, 38 p. Rocky Mountain Forest and Range Experiment Station, Fort Collins, Colo.
- Hoffman, George R., and Robert R. Alexander. 1980. Forest Vegetation of the Routt National Forest, Colorado: A habitat type classification. USDA Forest Service Research Paper RM-221, 41 p. Rocky Mountain Forest and Range Experiment Station, Fort Collins, Colo.
- Hoffman, George R. and Robert R. Alexander. 1983. Forest vegetation of the White River National Forest in western Colorado: A habitat type classification. USDA Forest Service Research Paper RM-249, 36 p. Rocky Mountain Forest and Range Experiment Station, Fort Collins, Colo.
- Johnson, James R., and James T. Nichols. 1970. Plants of South Dakota grasslands. A photographic study. South Dakota State University Agriculture Experiment Station Bulletin 566, 163 p. Brookings.
- Laird, Wilson M. 1950. The geology of the south unit, Theodore Roosevelt National Memorial Park. *North Dakota History* 17:225-240.
- Laird, Wilson M. 1956. The geology of the north unit, Theodore Roosevelt National Memorial Park. *North Dakota History* 23:53-77.
- Lauenroth, W. K., and W. C. Whitman. 1977. Dynamics of dry matter production in a mixed-grass prairie in western North Dakota. *Oecologia* 27:339-351.
- Layser, Earle F. 1974. Vegetative classification: Its application to forestry in the Northern Rocky Mountains. *Journal of Forestry* 72:354-357.
- Mackie, Richard J. 1970. Range ecology and relations of mule deer, elk and cattle in the Missouri River breaks, Montana. *Wildlife Monographs* 20, 79 p. Wildlife Society, Bethesda, Md.
- Moir, William H. 1972. Natural areas. *Science* 177:396-400.
- Moodie, C. D., and F. E. Koehler. 1975. Laboratory manual for soil fertility. 206 p. Student Book Corporation, Pullman, Wash.
- Mueggler, W. F., and W. L. Stewart. 1980. Grassland and shrubland habitat types of western Montana. USDA Forest Service General Technical Report INT-66, 154 p. Intermountain Forest and Range Experiment Station, Ogden, Utah.

- Nelson, Jack R. 1961. Composition and structure of the principal woody vegetation types in the North Dakota badlands. M.S. thesis, 195 p. North Dakota State University, Fargo.
- Omodt, H. W., G. A. Johnsgard, D. D. Patterson, and O. P. Olson. 1968. The major soils of North Dakota. North Dakota State University Agricultural Experiment Station Bulletin 472, 60 p. Fargo.
- Pfister, Robert D. 1972. Habitat types and regeneration. p. 120-125. In Permanent Association Committee proceedings. Western Forest and Conservation Association, Portland, Oreg.
- Pfister, Robert D., Bernard L. Kovalchik, Stephen F. Arno, and Richard C. Presley. 1977. Forest habitat types of Montana. USDA Forest Service General Technical Report INT-34, 175 p. Intermountain Forest and Range Experiment Station, Ogden, Utah.
- Quinnild, Clayton L., and Hugh E. Cosby. 1958. Relicts of climax vegetation on two mesas in western North Dakota. *Ecology* 39:29-32.
- Ralston, R. C. 1960. The structure and ecology of the north slope juniper stands of the Little Missouri Badlands. M.S. thesis, 85 p. University of Utah, Salt Lake City.
- Redmann, Robert E. 1975. Production ecology of grassland plant communities in western North Dakota. *Ecological Monographs* 45:86-106.
- Rickard, William H. 1957. The distribution of small mammals in relation to the climax vegetation mosaic in eastern Washington and northern Idaho. *Ecology* 41:99-106.
- Severson, Kieth E., and Charles E. Boldt. 1978. Cattle, wildlife, and riparian habitats in the western Dakotas. p. 90-103. In Regional rangeland symposium, Bismarck, N. Dak.
- Van Bruggen, Theodore. 1976. The vascular plants of South Dakota. 538 p. Iowa State University Press, Ames.
- Vaubel, John Arthur. 1975. Vegetation development in relation to age of river stabilization structures along a channelized segment of the Missouri River. M.A. thesis, 108 p. University of South Dakota, Vermillion.
- Weaver, J. E., and F. W. Albertson. 1956. Grasslands of the Great Plains. 395 p. Johnson Publishing Company, Lincoln, Nebr.
- Whitman, W., and H. C. Hanson. 1939. Vegetation on scoria and clay buttes in western North Dakota. *Ecology* 20:455-457.
- Wilson, Roger E. 1970. Succession in stands of *Populus deltoides* along the Missouri River in southeastern South Dakota. *American Midland Naturalist* 83:330-342.
- Wright, John C., and Elnora A. Wright. 1948. Grassland types of south central Montana. *Ecology* 29:449-460.



## **APPENDIX 1. HABITAT TYPE TABLES WITH STAND DATA**

In the plant data, the number to the left of the dot is percent coverage where the value exceeds 0.5%, with a + to the left of dot indicating coverage of 0.5% or less. Number to the right of dot is percent frequency. Species present in the macroplot but not in the microplots are indicated by an asterisk (\*). Stand numbers, locations, and topographic positions are also given.

Table A1.—*Stipa comata*/*Carex filifolia* habitat type

	Stand number						
	27	35	36	39	40	56	61
<b>Location</b>							
Quarter section	NW	SW	SE	NW	NE	SW	NW
Section	16	2	3	11	11	24	32
Township	140N	140N	140N	140N	140N	148N	148N
Range	102W	101W	101W	101W	101W	100W	100W
<b>Topographic position</b>							
Slope (%)	0	0	0	4	8	18	3
Aspect (°)	0	0	0	270	292	178	269
Elevation (m)	823	823	835	823	826	750	774
<b>Coverage/Frequency</b>							
<b>SHRUBS</b>							
<i>Artemisia cana</i>	.	.	+.2	.	.	.	.
<i>Artemisia frigida</i>	1.4	4.52	3.34	1.12	3.40	5.60	6.60
<i>Yucca glauca</i>	.	.	.	.	+.2	.	.
<b>GRAMINOIDS</b>							
<i>Agropyron caninum</i>	.	.	.	+.2	.	.	+.6
<i>Agropyron smithii</i>	68.98	1.12	+.8	1.18	2.6	.	.
<i>Andropogon scoparius</i>	.	1.4	.	.	.	.	.
<i>Aristida longiseta</i>	+.2	.	.	.	.	.	.
<i>Bouteloua gracilis</i>	.	1.20	2.24	+.2	2.18	+.8	1.6
<i>Bromus tectorum</i>	.	.	1.4	.	.	.	.
<i>Calamovilfa longifolia</i>	.	.	.	+.2	.	1.4	.
<i>Carex eleocharis</i>	4.26	1.22	1.28	2.28	1.14	+.8	+.6
<i>Carex filifolia</i>	48.99	24.84	37.98	15.80	22.76	49.99	4.36
<i>Koeleria pyramidata</i>	3.28	10.72	13.84	5.58	14.66	8.38	15.94
<i>Poa interior</i>	.	+.8	+.8	.	.	.	+.2
<i>Stipa comata</i>	11.64	41.94	52.99	65.99	35.88	35.98	35.98
<b>FORBS</b>							
<i>Artemisia campestris</i>	.	+.2	.	.	.	.	+.6
<i>Artemisia dracunculus</i>	5.22	4.34	1.28	+.4	5.42	3.32	1.6
<i>Aster oblongifolius</i>	.	.	.	.	.	+.2	.
<i>Astragalus crassicaupus</i>	.	.	.	+.2	.	.	.
<i>Astragalus lotiflorus</i>	.	.	+.2	.	.	.	.
<i>Chenopodium album</i>	.	.	+.2	.	.	.	.
<i>Coryphantha vivipara</i>	.	.	.	.	+.2	.	.
<i>Chrysopsis villosa</i>	.	6.32	1.14	2.4	2.24	.	14.64
<i>Echinacea angustifolia</i>	.	.	.	.	.	.	+.2
<i>Eriogonum flavum</i>	.	.	.	.	.	.	+.2
<i>Erysimum asperum</i>	.	+.2	+.2	.	.	.	.
<i>Gaura coccinea</i>	2.26	+.2	.	1.4	1.6	1.6	+.8
<i>Gentianella amarella</i>	.	.	.	.	.	.	+.2
<i>Helianthus rigidus</i>	.	.	.	.	.	1.6	.
<i>Lactuca oblongifolia</i>	1.20	6.78	3.42	2.44	8.82	8.76	.
<i>Liatris punctata</i>	.	+.8	+.10	+.8	.	+.2	1.12
<i>Linum perenne</i>	.	.	.	+.8	.	.	.
<i>Linum rigidum</i>	.	.	.	.	.	+.8	.
<i>Lithospermum incisum</i>	.	1.36	1.36	1.40	1.30	+.2	.
<i>Lygodesmia juncea</i>	.	2.40	1.40	+.18	1.32	2.24	.
<i>Melilotus officinalis</i>	.	+.4	.	.	.	.	.
<i>Opuntia polyacantha</i>	.	+.2	.	1.14	+.2	+.2	.
<i>Oxytropis lambertii</i>	1.10	+.2	.	.	.	.	7.70
<i>Petalostemon purpureum</i>	.	.	+.2	.	.	.	+.6
<i>Polygala alba</i>	.	.	.	.	.	+.6	+.2
<i>Potentilla pensylvanica</i>	.	+.2	.	.	+.4	+.4	.
<i>Psoralea argophylla</i>	4.44	+.6	.	+.2	+.18	+.8	.
<i>Ratibida columnifera</i>	1.8	+.2	.	.	.	+.2	.
<i>Selaginella densa</i>	39.92	43.92	43.99	54.90	42.88	28.96	84.99
<i>Sphaeralcea coccinea</i>	+.2	.	.	.	.	.	.
<i>Thermopsis rhombifolia</i>	.	.	.	.	+.4	.	.
<i>Tragopogon dubius</i>	+.12	+.6	+.4	+.8	+.8	1.10	+.2
<i>Vicia americana</i>	.	.	.	+.16	.	.	.
<i>Zigadenus venenosus</i>	.	+.8	+.8	+.8	+.6	+.6	+.2
<b>Summary</b>							
Species in microplots	16	27	23	24	22	24	22
Coverage of shrubs	1	4	3	1	3	5	6
Coverage of graminoids	134	79	106	88	76	96	55
Coverage of forbs	53	62	50	61	60	44	107
Total coverage	188	145	159	150	139	145	168



Table A2.—*Agropyron smithii*/*Carex filifolia* habitat type

	Stand number								
	3	5	14	18	20	21	49	54	62
<b>Location</b>									
Quarter section	SE	SE	SE	SE	SE	NW	NW	NE	NE
Section	1	1	12	28	28	34	33	26	32
Township	140N	140N	140N	141N	141N	141N	148N	148N	148N
Range	102W	102W	102W	101W	101W	101W	99W	100W	99W
<b>Topographic position</b>									
Slope (%)	16	19	16	18	18	20	14	9	14
Aspect (°)	203	202	43	233	137	211	163	357	124
Elevation (m)	701	719	707	732	713	719	631	750	643
<b>Coverage/Frequency</b>									
<b>SHRUBS</b>									
<i>Artemisia cana</i>	.	.	.	.	.	.	+.4	.	.
<i>Artemisia frigida</i>	+.4	+.6	.	2.20	.	3.26	1.10	3.18	1.4
<i>Atriplex nuttallii</i>	+.2	.	.	.	.	.	.	.	.
<i>Ceratoides lanata</i>	+.4	2.6	.	.	1.4	.	.	.	.
<b>GRAMINOIDS</b>									
<i>Agropyron smithii</i>	93.99	92.99	98.99	86.99	91.99	82.99	93.99	96.99	91.99
<i>Aristida longiseta</i>	.	.	.	.	.	1.4	.	.	.
<i>Bouteloua gracilis</i>	.	.	.	.	.	10.42	2.24	.	16.36
<i>Carex eleocharis</i>	.	.	.	.	.	.	.	1.4	.
<i>Carex filifolia</i>	28.62	.	.	27.72	29.84	17.64	1.4	24.94	+.2
<i>Koeleria pyramidata</i>	.	.	.	+.2	.	.	.	3.30	+.2
<i>Stipa comata</i>	.	.	.	3.16	4.34	2.16	1.20	+.2	+.2
<i>Stipa viridula</i>	3.12	4.14	.	3.14	.	.	.	.	4.22
<b>FORBS</b>									
<i>Achillea millefolium</i>	.	.	.	.	.	.	.	+.8	.
<i>Arabis holboellii</i>	.	.	.	.	.	+.4	.	.	.
<i>Artemisia dracuncululus</i>	.	.	.	.	.	.	.	+.2	.
<i>Conyza canadensis</i>	.	.	.	.	.	.	6.44	.	.
<i>Coryphantha vivipara</i>	+.2	.	.	.	.	.	+.2	.	.
<i>Euphorbia podperae</i>	+.2	.	.	.	.	.	.	.	.
<i>Gaura coccinea</i>	.	.	.	.	.	.	+.6	.	.
<i>Hedeoma hispida</i>	+.2	.	.	.	.	.	.	.	.
<i>Lactuca oblongifolia</i>	1.10	+.8	.	1.22	.	.	1.4	+.4	.
<i>Linum perenne</i>	.	.	+.2	+.2	.	.	.	.	.
<i>Melilotus officinalis</i>	.	.	.	.	.	.	.	.	+.6
Mosses & Lichens	2.28	+.2	.	+.2	.	.	.	1.4	.
<i>Opuntia polyacantha</i>	.	.	.	.	.	.	+.2	.	.
<i>Plantago patagonica</i>	.	.	.	.	.	.	2.32	.	.
<i>Polygala alba</i>	.	.	.	.	1.14	2.14	.	.	.
<i>Ratibida columnifera</i>	.	.	.	.	1.12	1.16	.	+.10	.
<i>Selaginella densa</i>	.	.	.	.	+.6	.	.	.	.
<i>Senecio plattensis</i>	.	.	.	.	+.4	+.4	.	.	.
<i>Sphaeralcea coccinea</i>	+.6	2.34	.	.	.	+.6	.	.	.
<i>Taraxacum officinale</i>	.	.	.	.	.	.	.	.	+.2
<i>Tragopogon dubius</i>	.	+.8	.	+.2	+.4	+.6	+.2	+.2	+.4
Species in microplots <sup>1</sup>	12	8	2	10	9	12	13	12	10
Coverage of shrubs	0	2	0	2	1	3	1	3	1
Coverage of graminoids	124	96	98	119	124	112	97	124	111
Coverage of forbs	3	2	0	1	2	3	9	1	0
Total coverage	127	100	98	122	127	118	107	128	112

<sup>1</sup>Mosses and lichens are counted as one species in the total list.

Table A3.—*Andropogon scoparius*/*Carex filifolia* habitat type

	Stand number								
	8	15	22	23	24	46	55	63	64
<b>Location</b>									
Quarter section	SE	NE	SE	SE	NW	SE	NE	NE	NE
Section	28	13	35	35	4	29	26	28	28
Township	141N	140N	141N	141N	140N	147N	148N	148N	148N
Range	101W	102W	101W	101W	101W	100W	100W	100W	100W
<b>Topographic position</b>									
Slope (%)	27	18	28	30	42	13	6	16	0
Aspect (°)	271	242	70	205	195	300	193	179	0
Elevation (m)	738	738	747	747	747	725	744	716	719
<b>Coverage/Frequency</b>									
<b>SHRUBS</b>									
<i>Artemisia cana</i>	.	.	.	3.14	.	.	.	.	.
<i>Artemisia frigida</i>	+ .8	2.28	+ .2	+ .4	+ .4	+ .2	+ .2	.	+ .10
<i>Ceratoides lanata</i>	.	.	.	.	+ .2	.	.	.	.
<i>Gutierrezia sarothrae</i>	.	.	+ .2	.	.	.	.	.	.
<i>Juniperus horizontalis</i>	1.4	.	.	.	.	+ .2	.	.	.
<i>Rhus aromatica</i>	.	.	1.4	.	.	.	.	.	.
<i>Rosa arkansana</i>	+ .10	+ .2	+ .6	.	.	1.16	.	.	+ .6
<i>Shepherdia argentea</i>	.	.	.	.	.	.	.	.	+ .2
<i>Symphoricarpos</i>	.	.	.	.	.	.	.	.	.
<i>occidentalis</i>	.	.	3.2	.	.	.	.	.	.
<i>Yucca glauca</i>	.	+ .2	.	.	.	.	+ .12	.	.
<b>GRAMINOIDS</b>									
<i>Agropyron smithii</i>	.	.	+ .4	+ .8	.	.	+ .2	.	.
<i>Andropogon scoparius</i>	68.99	88.99	88.99	72.98	65.99	84.99	70.98	77.99	81.99
<i>Bouteloua curtipendula</i>	.	.	3.16	14.44	19.80	.	+ .2	3.12	.
<i>Bouteloua gracilis</i>	.	.	.	.	1.10	.	+ .4	+ .2	.
<i>Calamovilfa longifolia</i>	6.30	+ .4	4.32	14.42	8.34	+ .2	14.74	+ .4	1.12
<i>Carex filifolia</i>	8.40	9.58	2.18	13.54	9.48	12.56	10.52	8.60	10.72
<i>Carex sprengei</i>	4.44	.	.	.	.	.	.	.	.
<i>Koeleria pyramidata</i>	+ .2	+ .8	1.18	.	+ .2	1.10	2.22	1.10	1.14
<i>Muhlenbergia cuspidata</i>	.	.	.	.	.	+ .2	1.8	.	.
<i>Stipa comata</i>	+ .2	.	+ .2	3.18	1.6	.	.	+ .4	.
<b>FORBS</b>									
<i>Agoseris glauca</i>	.	.	+ .2	.	.	.	.	.	.
<i>Anemone cylindrica</i>	.	.	.	.	.	+ .6	.	.	+ .2
<i>Anemone patens</i>	1.18	+ .4	.	.	.	.	.	.	.
<i>Antennaria parviflora</i>	+ .4	1.20	+ .2	.	.	.	.	.	.
<i>Antennaria rosea</i>	+ .4	.	.	.	.	+ .2	.	.	.
<i>Arabis holboellii</i>	.	+ .2	.	.	.	.	.	.	+ .2
<i>Artemisia dracunculus</i>	.	+ .4	.	.	.	.	.	.	.
<i>Aster laevis</i>	.	.	.	.	.	+ .4	.	.	.
<i>Aster oblongifolius</i>	1.26	1.8	+ .4	.	+ .4	+ .8	+ .8	+ .12	+ .6
<i>Astragalus adsurgens</i>	.	.	+ .10	.	.	.	.	.	.
<i>Astragalus crassicaulis</i>	.	+ .2	.	.	.	.	.	.	.
<i>Astragalus gilviflorus</i>	.	.	.	.	.	+ .4	.	.	.
<i>Calylophus serrulatus</i>	+ .4	+ .2	+ .2	.	.	.	.	.	.
<i>Campanula rotundifolia</i>	+ .6	+ .2	+ .2	.	.	+ .2	.	.	.
<i>Castilleja sessiliflora</i>	.	.	+ .2	.	.	.	.	.	.
<i>Chrysopsis villosa</i>	.	+ .2	.	.	.	.	.	.	.
<i>Cirsium undulatum</i>	.	.	.	.	1.4	.	.	.	.
<i>Echinacea angustifolia</i>	2.34	+ .6	1.12	.	+ .8	+ .8	+ .14	1.26	+ .12
<i>Erigeron strigosus</i>	.	.	+ .2	.	.	+ .10	+ .2	.	.
<i>Galium boreale</i>	.	.	+ .2	.	.	.	.	.	.
<i>Gaura coccinea</i>	1.14	+ .4	+ .2	+ .4	.	.	2.36	1.22	.
<i>Glycyrrhiza lepidota</i>	.	.	.	.	.	.	.	1.14	.
<i>Hedeoma hispida</i>	.	.	.	.	1.10	.	.	.	.
<i>Helianthus rigidus</i>	3.20	+ .4	7.52	.	.	7.66	7.86	2.56	6.70
<i>Lactuca oblongifolia</i>	2.26	+ .12	.	2.36	4.58	1.30	.	+ .2	.
<i>Liatis punctata</i>	.	.	.	.	.	+ .2	.	.	.
<i>Linum perenne</i>	+ .6	+ .2	+ .6	.	.	.	.	.	.
<i>Linum rigidum</i>	.	.	.	.	+ .2	.	.	.	.
<i>Lithospermum incisum</i>	.	.	.	.	.	.	.	.	+ .2
<i>Lygodesmia juncea</i>	+ .10	.	+ .2	.	+ .8	+ .4	+ .8	.	1.4
<i>Medicago sativa</i>	.	.	.	.	.	.	.	.	+ .2
<i>Melilotus officinalis</i>	.	.	.	.	.	+ .2	.	.	1.16
Mosses & Lichens	2.38	.	.	.	.	.	.	.	.
<i>Opuntia fragilis</i>	.	.	.	+ .2	.	.	.	.	.
<i>Opuntia polyacantha</i>	.	+ .4	+ .4	.	.	.	.	.	.
<i>Oxytropis lambertii</i>	.	.	.	.	+ .4	.	+ .4	+ .4	+ .2
<i>Petalostemon purpureum</i>	+ .6	.	+ .2	.	.	.	.	.	+ .2
<i>Polygala alba</i>	+ .4	.	+ .4	+ .2	.	+ .2	+ .4	.	.
<i>Potentilla arguta</i>	.	.	+ .4	+ .2	.	.	.	.	.
<i>Psoralea argophylla</i>	.	.	.	2.16	2.34	.	1.48	.	+ .8
<i>Psoralea esculenta</i>	.	+ .2	.	.	.	.	+ .2	+ .2	.
<i>Senecio plattensis</i>	+ .16	+ .6	.	1.10	+ .2	+ .2	.	.	.
<i>Solidago rigida</i>	1.12	.	.	.	.	.	.	.	.
<i>Sphaeralcea coccinea</i>	.	.	+ .8	+ .4	1.10	.	.	.	.
<i>Tragopogon dubius</i>	.	+ .2	.	.	+ .2	.	.	.	.
<i>Vicia americana</i>	.	.	.	.	.	.	.	+ .8	1.4
<b>Summary</b>									
Species in microplots	26	25	32	15	22	23	20	16	20
Coverage of shrubs	1	2	4	3	0	1	0	0	0
Coverage of graminoids	86	97	98	116	103	97	97	89	94
Coverage of forbs	13	2	9	5	9	8	10	5	9
Total coverage	100	101	111	124	112	106	107	94	103



Table A4.—*Juniperus horizontalis*/*Andropogon scoparius* habitat type

	Stand number				
	9	10	12	44	47
<b>Location</b>					
Quarter section	NE	SE	SW	NW	SE
Section	1	15	12	22	33
Township	140N	140N	140N	140N	148N
Range	102W	102W	102W	101W	99W
<b>Topographic position</b>					
Slope (%)	28	48	30	46	32
Aspect (°)	349	268	9	21	338
Elevation (m)	719	725	719	753	628
<b>Coverage/Frequency</b>					
<b>SHRUBS</b>					
<i>Artemisia cana</i>	.	3.4	.	.	.
<i>Artemisia frigida</i>	.	.	+ .2	.	.
<i>Juniperus communis</i>	+ .8	.	.	2.2	.
<i>Juniperus horizontalis</i>	61.92	82.99	85.99	85.98	90.99
<i>Potentilla fruticosa</i>	1.2	7.12	3.16	1.10	.
<i>Prunus virginiana</i>	.	2.8	.	.	.
<i>Rhus aromatica</i>	+ .4	7.16	1.2	.	.
<i>Rosa arkansana</i>	+ .2	+ .8	+ .2	.	.
<i>Shepherdia argentea</i>	.	1.2	.	.	.
<i>Symphoricarpos occidentalis</i>	.	7.60	2.36	+ .4	.
<b>GRAMINOIDS</b>					
<i>Agropyron caninum</i>	.	.	.	+ .4	+ .2
<i>Andropogon scoparius</i>	36.92	25.80	24.84	35.84	2.6
<i>Aristida longiseta</i>	.	.	.	+ .2	.
<i>Bouteloua curtipendula</i>	.	.	.	+ .4	.
<i>Calamovilfa longifolia</i>	8.54	4.32	15.48	8.36	2.12
<i>Carex filifolia</i>	1.16	1.12	5.20	5.36	3.30
<i>Carex sprengei</i>	+ .6	1.16	.	.	.
<i>Koeleria pyramidata</i>	1.4	+ .6	+ .4	+ .4	1.4
<i>Muhlenbergia cuspidata</i>	.	.	.	3.12	.
<i>Stipa spartea</i>	.	.	.	1.10	31.88
<b>FORBS</b>					
<i>Achillea millefolium</i>	.	+ .12	+ .2	.	.
<i>Anemone cylindrica</i>	.	.	.	1.42	.
<i>Anemone patens</i>	4.54	3.24	3.50	.	2.52
<i>Antennaria parviflora</i>	+ .2	.	+ .2	.	.
<i>Antennaria rosea</i>	.	.	.	+ .2	+ .2
<i>Artemisia campestris</i>	+ .6	.	.	.	.
<i>Aster falcatus</i>	.	2.12	.	.	.
<i>Aster laevis</i>	+ .2	.	.	+ .16	2.28
<i>Aster oblongifolius</i>	1.12	.	+ .2	.	+ .4
<i>Astragalus adsurgens</i>	+ .2	.	.	.	.
<i>Astragalus gilviflorus</i>	+ .6	1.14	2.26	.	.
<i>Astragalus</i> sp.	.	.	1.4	.	.
<i>Campanula rotundifolia</i>	2.18	1.14	1.12	+ .14	2.32
<i>Chenopodium album</i>	.	.	.	+ .2	.
<i>Echinacea angustifolia</i>	+ .14	1.14	1.18	.	+ .2
<i>Erigeron strigosus</i>	.	.	.	.	+ .8
<i>Eriogonum flavum</i>	+ .4	.	.	.	.
<i>Galium boreale</i>	.	9.66	6.76	3.72	6.82
<i>Gaura coccinea</i>	1.6	1.10	+ .14	.	1.22
<i>Geum triflorum</i>	.	.	.	.	+ .2
<i>Hedysarum boreale</i>	12.40	.	.	.	.
<i>Helianthus rigidus</i>	2.24	+ .4	+ .8	.	.
<i>Liatris punctata</i>	+ .10	+ .2	.	.	.
<i>Linum perenne</i>	+ .8	+ .8	.	+ .4	+ .6
<i>Linum rigidum</i>	+ .2	.	.	.	.
<i>Lygodesmia juncea</i>	1.10	.	+ .8	.	.
<i>Melilotus albus</i>	.	.	.	.	+ .2
<i>Melilotus officinalis</i>	.	.	.	1.4	4.26
Mosses & Lichens	+ .4	1.10	1.10	.	.
<i>Opuntia polyacantha</i>	.	.	.	+ .2	.
<i>Oxytropis lambertii</i>	+ .4	.	.	+ .2	+ .8
<i>Petalostemon purpureum</i>	1.18	+ .2	+ .14	1.12	1.22
<i>Polygala alba</i>	+ .4	.	+ .2	.	.
<i>Potentilla pensylvanica</i>	.	.	1.2	.	.
<i>Psoralea esculenta</i>	+ .4	.	+ .2	.	.
<i>Senecio plattensis</i>	+ .8	+ .6	+ .4	1.10	.
<i>Solidago nemoralis</i>	+ .8	.	.	.	.
<i>Solidago rigida</i>	+ .2	.	.	.	.
<i>Thermopsis rhombifolia</i>	1.18	+ .2	.	.	.
<i>Vicia americana</i>	.	2.22	+ .6	.	.
<i>Viola</i> sp.	.	.	.	.	+ .2
<b>Summary</b>					
Species in microplots	36	29	29	25	23
Coverage of shrubs	62	109	91	88	90
Coverage of graminoids	46	31	44	52	39
Coverage of forbs	25	21	16	7	18
Total coverage	133	161	151	147	147

Table A5.—*Artemisia tridentata*/*Agropyron smithii* habitat type

	Stand number			
	4	13	16	28
<b>Location</b>				
Quarter section	SE	SW	SW	NW
Section	1	12	20	4
Township	140N	140N	140N	140N
Range	102W	102W	101W	101W
<b>Topographic position</b>				
Slope (%)	49	5	10	22
Aspect (°)	77	185	97	45
Elevation (m)	725	732	783	765
<b>Coverage/Frequency</b>				
<b>SHRUBS</b>				
<i>Artemisia frigida</i>	.	.	+ .2	+ .2
<i>Artemisia tridentata</i>	17.56	36.88	27.72	41.82
<i>Atriplex confertifolia</i>	4.14	.	2.2	+ .2
<i>Atriplex nuttallii</i>	.	+ .2	.	.
<i>Ceratoides lanata</i>	+ .2	.	2.4	+ .2
<i>Eriogonum pauciflorum</i>	.	.	.	+ .2
<i>Gutierrezia sarothrae</i>	.	1.8	.	+ .2
<i>Ribes setosum</i>	.	.	+ .4	.
<i>Symphoricarpos occidentalis</i>	.	.	4.16	.
<b>GRAMINOIDS</b>				
<i>Agropyron caninum</i>	7.28	20.50	4.16	2.6
<i>Agropyron smithii</i>	64.99	45.90	86.98	81.99
<i>Andropogon scoparius</i>	+ .2	.	.	.
<i>Bouteloua gracilis</i>	.	.	.	+ .2
<i>Carex filifolia</i>	.	.	1.16	12.90
<i>Muhlenbergia cuspidata</i>	.	.	3.24	+ .2
<i>Koeleria pyramidata</i>	.	.	.	3.6
<i>Stipa viridula</i>	.	.	1.4	3.14
<b>FORBS</b>				
<i>Achillea millefolium</i>	.	.	4.18	1.6
<i>Astragalus agrestis</i>	1.8	+ .6	.	.
<i>Gaura coccinea</i>	.	.	.	+ .2
<i>Geum triflorum</i>	.	+ .4	.	.
<i>Lactuca oblongifolia</i>	+ .6	.	2.30	+ .4
Mosses & Lichens	+ .2	.	.	.
<i>Opuntia polyacantha</i>	+ .2	+ .4	+ .4	1.8
<i>Parietaria pennsylvanica</i>	.	.	2.10	.
<i>Polygala alba</i>	.	.	.	1.4
<i>Ratibida columnifera</i>	.	.	.	1.14
<i>Sphaeralcea coccinea</i>	+ .2	+ .2	.	.
<i>Vicia americana</i>	1.14	+ .4	+ .2	.
<b>Summary</b>				
Species in microplots	12	10	16	19
Coverage of shrubs	21	37	35	41
Coverage of graminoids	71	65	95	101
Coverage of forbs	2	0	8	4
Total coverage	94	102	138	146



Table A6.—*Artemisia cana*/Agropyron smithii habitat type

	Stand number											
	1	6	7	26	66	69	70	2	25	31	32	65
Location												
Quarter section	SE	NW	NW	SE	NW	NW	NW	SE	NW	NE	SE	SW
Section	1	6	6	35	30	30	30	1	4	14	1	33
Township	140N	140N	140N	140N	148N	148N	148N	140N	140N	140N	140N	148N
Range	102W	101W	101W	101W	99W	99W	99W	102W	101W	102W	102W	99W
Topographic position												
Slope (%)	0	5	5	5	0	0	4	0	8	7	0	0
Aspect (°)	0	245	287	32	0	0	76	0	293	245	0	0
Elevation (m)	683	707	706	744	613	619	619	683	744	683	689	600
Coverage/Frequency												
SHRUBS												
<i>Artemisia cana</i>	24.80	48.88	26.70	36.74	34.62	26.50	42.80	32.64	45.82	18.48	51.96	21.66
<i>Artemisia frigida</i>	.	+ 2	+ 2	+ 4	.	.	.	.	3.6	+ 6	1.8	3.16
<i>Ceratoides lanata</i>	.	4.14	1.2	+ 4	.	.	.	.	2.8	.	.	.
<i>Gutierrezia sarothrae</i>	.	.	.	.	+ 2	.	.	.	.	.	.	+ 2
<i>Rosa woodsii</i>	.	.	.	.	.	.	.	.	.	.	.	1.2
<i>Symphoricarpos occidentalis</i>	.	.	.	.	.	.	.	63.82	24.48	61.72	13.26	70.88
GRAMINOIDS												
<i>Agropyron caninum</i>	.	+ 4	2.10	4.10	.	.	.	7.20	3.8	23.72	13.32	27.54
<i>Agropyron smithii</i>	79.99	80.99	88.99	73.94	95.99	96.99	97.99	31.52	84.99	25.42	57.93	46.76
<i>Bouteloua gracilis</i>	.	.	.	.	.	.	+ 8	.	.	.	.	.
<i>Bromus inermis</i>	.	.	.	.	.	.	.	.	.	.	1.2	.
<i>Calamagrostis inexpansa</i>	.	.	.	.	.	+ 2	1.6	.	.	.	.	.
<i>Calamovilfa longifolia</i>	.	.	.	5.10	.	.	.	.	.	2.4	.	.
<i>Carex filifolia</i>	.	.	.	+ 4	.	.	.	.	+ 2	.	.	.
<i>Distichlis spicata</i>	.	.	.	.	.	+ 2	.	.	5.10	.	.	.
<i>Muhlenbergia cuspidata</i>	.	.	.	5.12	.	.	.	.	.	.	.	.
<i>Muhlenbergia racemosa</i>	.	.	.	.	.	.	.	.	.	+ 2	.	.
<i>Poa arida</i>	.	.	.	+ 6	.	.	.	.	.	.	.	7.26
<i>Poa interior</i>	9.44	1.4	.	2.6	.	.	.	1.6	.	.	.	.
<i>Poa palustris</i>	.	.	.	1.8	.	.	.	1.4	4.18	.	.	.
<i>Poa pratensis</i>	.	.	.	.	.	.	.	.	.	13.32	.	2.4
<i>Stipa comata</i>	.	.	.	1.4	.	.	.	.	.	.	.	.
<i>Stipa viridula</i>	.	2.12	+ 4	11.30	.	3.16	2.12	.	1.6	1.8	+ 4	8.28
FORBS												
<i>Achillea millefolium</i>	7.36	3.14	.	+ 4	1.2	.	.	.	1.2	2.16	.	4.32
<i>Artemisia dracunculus</i>	.	.	.	.	.	.	.	.	.	.	.	+ 2
<i>Artemisia ludoviciana</i>	.	2.6	.	.	.	.	.	.	.	6.18	.	.
<i>Aster oblongifolius</i>	.	.	.	.	.	.	.	.	.	+ 4	.	.
<i>Astragalus agrestis</i>	.	.	.	.	.	.	.	.	1.8	.	.	.
<i>Chenopodium album</i>	.	.	.	.	.	.	.	+ 8	.	.	.	.
<i>Convolvulus sepium</i>	.	.	.	.	.	.	.	.	.	.	+ 2	.
<i>Conyza canadensis</i>	.	.	.	.	.	.	1.12	.	.	.	3.28	.
<i>Erigeron strigosus</i>	.	.	.	.	.	.	+ 4	.	.	.	.	.
<i>Euphorbia podperae</i>	.	+ 6	.	.	.	.	.	.	.	.	.	.
<i>Euphorbia spathulata</i>	.	.	.	.	.	.	.	.	+ 2	.	.	.
<i>Galium aparine</i>	.	+ 4	.	.	.	.	.	1.8	.	.	.	.
<i>Gaura coccinea</i>	.	.	.	2.14	.	.	.	.	1.6	1.4	+ 8	.
<i>Grindelia squarrosa</i>	.	.	.	.	+ 2	.	.	.	.	.	.	.
<i>Hedeoma hispida</i>	.	.	.	.	.	.	.	+ 2	+ 2	.	.	.
<i>Helianthus rigidus</i>	.	.	.	.	.	.	.	.	+ 2	.	.	.
<i>Lactuca oblongifolia</i>	.	.	.	11.38	+ 2	.	.	1.16	10.54	.	.	.
<i>Linum perenne</i>	.	.	.	+ 4	.	.	.	.	2.12	+ 2	.	.
<i>Melilotus officinalis</i>	.	.	.	+ 6	1.10	+ 2	.	.	.	+ 2	3.24	.
<i>Monarda fistulosa</i>	.	.	.	.	.	.	.	.	+ 2	.	.	.
Mosses & Lichens	11.70	.	.	.	.	.	.	+ 4	.	.	.	.
<i>Opuntia fragilis</i>	.	.	.	.	.	.	.	.	.	.	+ 2	.
<i>Opuntia polyacantha</i>	.	+ 2	.	.	.	.	.	.	.	.	.	.
<i>Parietaria pennsylvanica</i>	.	15.44	1.6	.	.	.	.	3.48	+ 6	.	2.12	.
<i>Plantago patagonica</i>	.	.	.	.	.	.	.	.	.	.	+ 4	.
<i>Psoralea esculenta</i>	.	.	.	6.24	.	.	.	.	.	.	.	.
<i>Ratibida columnifera</i>	.	.	.	+ 2	.	.	1.4	.	+ 6	.	.	.
<i>Solidago missouriensis</i>	.	.	.	.	.	.	.	.	.	.	2.4	.
<i>Solidago rigida</i>	.	.	.	.	.	.	.	.	1.2	.	.	.
<i>Sphaeralcea coccinea</i>	.	1.10	+ 2	.	.	.	.	.	.	+ 2	.	.
<i>Taraxacum officinale</i>	.	.	.	.	.	.	+ 4	.	.	.	.	.
<i>Thalictrum dasycarpum</i>	.	.	.	.	.	.	.	.	.	1.4	.	.
<i>Tragopogon dubius</i>	.	.	.	.	.	.	+ 2	.	.	.	.	.
<i>Vicia americana</i>	.	1.4	.	.	1.10	.	.	1.12	1.10	+ 2	1.12	.
Species in microplots	5	14	8	20	8	6	10	13	23	18	16	12
Coverage of shrubs	24	52	27	36	34	26	42	95	74	79	65	95
Coverage of graminoids	88	83	90	102	95	99	100	40	97	63	71	90
Coverage of forbs	18	22	1	19	3	0	2	6	17	10	11	4
Total coverage	130	157	118	157	132	125	144	141	188	152	147	189

Table A7.—*Fraxinus pennsylvanica*/*Symphoricarpos occidentalis*, *Fraxinus pennsylvanica*/*Prunus virginiana*, and *Populus tremuloides*/*Betula occidentalis* habitat types

	Stand number											
	<i>Fraxinus</i> / <i>Symphoricarpos</i> H.T.				<i>Fraxinus</i> / <i>Prunus</i> H.T.				<i>Populus</i> / <i>Betula</i> H.T.			
	29	30	51	52	41	42	57	58	43	60	67	68
Location												
Quarter section	SE	SE	NW	SW	SW	SW	SE	NW	SW	NE	NW	SE
Section	16	16	31	31	11	11	21	27	35	32	11	3
Township	140N	140N	148N	148N	140N	140N	148N	148N	140N	148N	147N	147N
Range	102W	102W	99W	99W	102W	102W	100W	100W	101W	100W	100W	100W
Topographic position												
Slope (%)	0	11	2	0	0	3	6	14	77	13	23	48
Aspect (°)	0	28	24	0	0	283	340	25	30	73	341	320
Elevation (m)	683	683	600	600	710	707	732	728	838	753	765	759
Coverage/Frequency												
SHRUBS												
<i>Amelanchier alnifolia</i>	.	.	.	.	.	.	.	.	.	4.4	.	.
<i>Arctostaphylos uva-ursi</i>	.	.	.	.	.	.	.	.	.	.	.	.
<i>Betula occidentalis</i>	.	.	.	.	.	.	.	.	7.30	1.6	32.68	49.74
<i>Celastrus scandens</i>	.	.	.	.	.	.	.	.	+ .2	1.8	+ .2	+ .2
<i>Clematis ligusticifolia</i>	+ .4	.	1.4	2.12	+ .2	1.10	.	+ .2	.	.	.	+ .4
<i>Cornus stolonifera</i>	.	.	+ .2	.	.	.	.	.	1.2	+ .2	.	2.4
<i>Juniperus communis</i>	1.12	+ .6	.	.	.	.	.	.	1.20	.	.	+ .2
<i>Juniperus horizontalis</i>	.	.	.	.	.	.	.	.	.	.	.	.
<i>Parthenocissus vitacea</i>	+ .4	+ .2	2.10	1.2	.	.	.	.	.	.	.	.
<i>Potentilla fruticosa</i>	.	.	.	.	.	.	.	.	.	.	.	.
<i>Prunus virginiana</i>	.	.	.	.	17.40	44.76	19.32	31.70	36.76	37.70	35.80	16.50
<i>Rhus aromatica</i>	11.32	3.8	1.2	8.18	+ .2	3.4	.	.	+ .2	.	.	.
<i>Ribes odoratum</i>	.	.	.	.	+ .2	+ .2	.	+ .2	.	.	.	.
<i>Ribes setosum</i>	1.12	1.2	.	.	.	.	3.16	+ .4	+ .2	1.10	1.2	+ .2
<i>Rosa woodsii</i>	1.8	2.10	9.56	13.50	2.8	16.50	.	.	+ .6	+ .6	3.22	.
<i>Rubus idaeus</i>	.	.	.	.	.	.	.	.	.	4.14	.	.
<i>Salix bebbiana</i>	.	.	.	.	.	.	.	.	.	.	.	.
<i>Shepherdia argentea</i>	+ .2	1.4	.	.	.	.	.	.	9.28	.	.	.
<i>Symphoricarpos albus</i>	.	.	.	.	.	.	.	.	.	.	31.86	10.40
<i>Symphoricarpos occidentalis</i>	22.90	14.50	45.94	81.99	18.76	23.84	1.4	7.36	21.80	6.46	.	.
<i>Toxicodendron rydbergii</i>	17.62	2.14	61.99	12.28	+ .2	.	.	.	30.82	6.46	21.76	+ .4
<i>Vitis vulpina</i>	.	.	.	.	+ .2	.	.	.	.	.	.	.
GRAMINOIDS												
<i>Agropyron caninum</i>	9.36	5.24	1.6	2.8	.	.	1.6	.	.	.	2.10	.
<i>Bromus inermis</i>	.	.	.	.	.	.	.	.	.	.	.	.
<i>Calamovilfa longifolia</i>	16.36	29.64	.	.	.	.	.	.	.	.	.	.
<i>Carex sprengeii</i>	.	+ .2	.	.	.	.	52.88	70.92	.	25.44	9.18	.
<i>Elymus canadensis</i>	12.38	25.66	7.22	2.12	.	.	.	.	+ .2	.	.	.
<i>Elymus villosus</i>	.	.	.	.	.	.	.	29.66	.	.	.	.
<i>Elymus virginicus</i>	.	.	.	.	38.96	41.90	30.90	1.2	1.6	.	.	.
<i>Muhlenbergia racemosa</i>	8.4	5.22	15.64	5.14	.	.	.	1.6	.	+ .2	.	.
<i>Oryzopsis micrantha</i>	.	.	.	2.6	.	.	+ .2	.	36.86	10.30	.	.
<i>Poa paulstris</i>	.	.	.	.	.	.	1.6	.	.	.	.	.
<i>Poa pratensis</i>	60.90	51.94	1.6	.	6.10	17.26	39.60	.	+ .2	.	.	.
<i>Schizachne purpurascens</i>	.	.	.	.	.	.	3.10	.	.	.	.	.
<i>Scirpus americanus</i>	.	.	1.4	.	.	.	.	.	.	.	.	.
<i>Stipa viridula</i>	.	1.4	.	.	.	.	+ .2	.	.	.	.	.
FORBS												
<i>Achillea millefolium</i>	.	.	.	.	.	.	+ .6	.	1.12	+ .2	.	.
<i>Ambrosia psilostachya</i>	.	1.10	2.12	.	.	.	.	.	.	.	.	.
<i>Anemone cylindrica</i>	.	.	+ .2	.	.	.	.	1.2	+ .6	.	.	.
<i>Antennaria plantaginifolia</i>	.	1.8	.	.	.	.	.	.	.	.	.	.
<i>Apocynum androsaemifolium</i>	.	.	.	.	.	.	.	.	.	33.72	12.42	1.4
<i>Apocynum cannabinum</i>	.	.	.	.	.	.	.	.	.	4.22	.	.
<i>Aralia nudicaulis</i>	.	.	.	.	.	.	.	.	.	.	20.64	.
<i>Arctium minus</i>	.	.	.	.	9.64	1.14	+ .2	.	.	.	.	.



Table A7.—*Fraxinus pennsylvanica*/*Symphoricarpos occidentalis*, *Fraxinus pennsylvanica*/*Prunus virginiana*, and *Populus tremuloides*/*Betula occidentalis* habitat types—Continued

	Stand number											
	<i>Fraxinus</i> / <i>Symphoricarpos</i> H.T.				<i>Fraxinus</i> / <i>Prunus</i> H.T.				<i>Populus</i> / <i>Betula</i> H.T.			
	29	30	51	52	41	42	57	58	43	60	67	68
Location												
Quarter section	SE	SE	NW	SW	SW	SW	SE	NW	SW	NE	NW	SE
Section	16	16	31	31	11	11	21	27	35	32	11	3
Township	140N	140N	148N	148N	140N	140N	148N	148N	140N	148N	147N	147N
Range	102W	102W	99W	99W	102W	102W	100W	100W	101W	100W	100W	100W
Topographic position												
Slope (%)	0	11	2	0	0	3	6	14	77	13	23	48
Aspect (°)	0	28	24	0	0	283	340	25	30	73	341	320
Elevation (m)	683	683	600	600	710	707	732	728	838	753	765	759
Coverage/Frequency												
<i>Artemisia ludoviciana</i>	+ .4	*	1.6	.	.	.	*	.	.	.	.	.
<i>Asclepias verticillata</i>	.	.	.	.	.	.	.	.	.	.	.	.
<i>Asparagus officinalis</i>	*	*	.	.	.	.	.	.	.	.	.	.
<i>Aster laevis</i>	.	.	.	.	.	.	+ .2	+ .8	+ .2	.	.	.
<i>Astragalus agrestis</i>	.	.	.	.	.	.	.	.	+ .4	.	.	.
<i>Campanula rotundifolia</i>	.	.	.	.	.	.	.	.	+ .4	.	+ .2	.
<i>Chenopodium album</i>	.	.	.	+ .6	+ .6	.	.	.	.	.	.	.
<i>Convolvulus arvensis</i>	.	.	.	+ .4	.	.	.	.	.	.	.	.
<i>Convolvulus sepium</i>	.	.	.	.	.	.	+ .4	.	.	.	.	.
<i>Disporum trachycarpum</i>	.	.	.	.	.	.	.	1.4	2.26	+ .2	4.32	15.86
<i>Euphorbia podperae</i>	+ .2	.	.	.	.	.	.	.	.	.	.	.
<i>Fragaria virginiana</i>	.	.	.	.	.	.	.	3.40	1.10	.	4.60	1.16
<i>Galium aparine</i>	.	.	.	.	.	.	.	1.4	.	.	.	.
<i>Galium boreale</i>	4.10	.	.	4.28	19.74	3.14	12.76	7.56	5.34	8.46	7.38	+ .2
<i>Geranium carolinianum</i>	.	.	.	.	.	.	.	.	.	1.4	.	.
<i>Glycyrrhiza lepidota</i>	.	.	2.10	.	.	.	.	.	.	.	.	.
<i>Hackelia deflexa</i>	.	.	.	+ .2	.	.	.	.	.	.	.	.
<i>Helianthus rigidus</i>	.	.	+ .4	.	.	.	.	.	.	.	.	.
<i>Heuchera richardsonii</i>	.	.	.	.	.	.	.	1.8	1.14	+ .2	.	.
<i>Lactuca oblongifolia</i>	.	.	1.12	+ .4	5.68	4.54	1.10	.	.	+ .2	.	.
<i>Linum perenne</i>	.	.	.	.	.	.	.	.	.	.	.	.
<i>Lygodesmia juncea</i>	.	2.10	.	.	.	.	.	.	.	.	.	.
<i>Medicago sativa</i>	.	.	.	.	.	.	.	.	.	.	.	.
<i>Melilotus albus</i>	3.18	23.52	1.4	1.4	.	.	.	.	.	.	.	.
<i>Melilotus officinalis</i>	25.86	55.92	26.90	9.46	+ .4	.	.	.	.	.	.	.
<i>Monarda fistulosa</i>	.	.	.	.	4.14	1.6	1.4	.	1.14	.	.	.
Mosses & Lichens	.	.	.	.	.	.	.	.	.	1.2	.	3.4
<i>Osmorhiza longistylis</i>	.	.	.	.	.	.	.	.	.	.	.	.
<i>Oxalis stricta</i>	.	.	.	.	1.10	1.16	1.8	+ .2	.	.	.	.
<i>Parietaria pennsylvanica</i>	.	.	.	.	1.16	4.42	+ .2	.	.	.	.	.
Polypodiaceae	.	.	.	.	.	.	*	+ .8	1.14	4.6	.	.
<i>Ranunculus abortivus</i>	.	.	.	.	2.18	+ .10	.	.	.	.	.	.
<i>Ratibida columnifera</i>	*	.	.	.	.	.	.	.	.	.	.	.
<i>Sanicula marilandica</i>	.	.	.	.	.	.	2.4	1.18	.	.	.	19.70
<i>Smilacina stellata</i>	+ .8	3.18	+ .8	1.10	.	.	1.10	2.30	7.46	6.56	2.30	+ .6
<i>Smilax herbacea</i>	.	.	.	.	.	.	+ .6	+ .2	.	+ .2	.	.
<i>Solidago missouriensis</i>	.	.	*	+ .4	.	.	.	.	.	.	.	.
<i>Solidago rigida</i>	.	.	.	.	.	.	.	.	+ .4	.	.	.
<i>Taraxacum officinale</i>	+ .2	.	+ .6	+ .4	1.26	1.26	2.22	+ .6	+ .4	1.10	.	.
<i>Thalictrum dasycarpum</i>	*	.	*	.	2.18	3.22	*	*	.	.	+ .2	.
<i>Thalictrum venulosum</i>	.	.	*	+ .2	.	.	1.4	+ .2	.	*	2.10	.
<i>Thermopsis rhombifolia</i>	.	.	+ .2	.	.	.	.	.	1.8	.	.	.
<i>Tragopogon dubius</i>	+ .2	.	.	.	.	.	2.10	+ .2	.	.	+ .2	.
<i>Trifolium pratense</i>	2.16	4.16	.	.	.	.	.	.	.	.	.	.
<i>Urtica dioica</i>	.	.	1.4	1.6	1.4	+ .6	1.8	1.14	.	1.10	4.24	.
<i>Vicia americana</i>	1.8	*	+ .4	.	.	.	.	.	.	.	.	.
<i>Viola pratensis</i>	+ .12	+ .4	+ .2	+ .4	.	.	.	.	.	.	.	.
<i>Viola</i> sp.	.	.	.	.	.	.	+ .2	.	+ .2	.	.	.
Species in microplots	25	26	26	20	24	19	28	25	31	27	20	16
Coverage of shrubs	53	23	119	112	37	87	23	38	106	60	123	77
Coverage of graminoids	105	116	25	11	44	58	126	101	37	35	11	0
Coverage of forbs	35	89	34	16	45	29	24	18	20	59	55	39
Total coverage	193	228	178	139	126	164	173	157	163	154	189	116

Table A8.—*Juniperus scopulorum*/*Oryzopsis micrantha* habitat type

	Stand number						
	33	34	37	38	49	50	53
Location							
Quarter section	SW	SW	NW	SE	SW	SW	NE
Section	11	11	24	23	30	30	25
Township	140N	140N	140N	140N	148N	148N	148N
Range	101W	101W	101W	101W	99W	99W	100W
Topographic position							
Slope (%)	45	65	42	38	47	67	64
Aspect (°)	332	2	296	11	304	294	358
Elevation (m)	817	792	786	826	634	655	652
	Coverage/Frequency						
SHRUBS							
<i>Artemisia cana</i>	.	.	.	.	.	*	.
<i>Artemisia frigida</i>	.	.	+ .4	.	.	.	.
<i>Artemisia tridentata</i>	+ .2	.	.	.	.	+ .2	+ .2
<i>Ceratoides lanata</i>	.	.	.	.	.	.	*
<i>Chrysothamnus nauseosus</i>	.	.	+ .2	.	.	.	.
<i>Clematis ligusticifolia</i>	.	.	.	.	.	.	.
<i>Gutierrezia sarothrae</i>	.	.	.	.	.	.	*
<i>Potentilla fruticosa</i>	+ .2	.	.	.	.	.	.
<i>Prunus virginiana</i>	48.50	.	14.44	4.22	6.22	+ .2	2.8
<i>Rhus aromatica</i>	11.36	+ .2	12.36	2.10	2.8	3.6	+ .6
<i>Ribes odoratum</i>	+ .2	2.4	.	.	.	.	.
<i>Ribes setosum</i>	.	1.2	*	.	.	.	.
<i>Rosa arkansana</i>	+ .4	.	.	.	.	.	.
<i>Rosa woodsii</i>	+ .6	.	1.6	1.4	*	.	.
<i>Symphoricarpos occidentalis</i>	+ .8	3.10	3.34	2.20	4.24	1.8	+ .12
GRAMINOIDS							
<i>Agropyron caninum</i>	+ .4	+ .2	+ .2	5.26	+ .2	+ .2	*
<i>Carex eburnea</i>	.	.	.	.	.	6.10	4.8
<i>Oryzopsis micrantha</i>	59.99	62.99	68.99	68.98	59.96	77.99	76.99
<i>Poa pratensis</i>	.	.	.	.	.	.	+ .2
<i>Stipa viridula</i>	.	.	.	.	.	.	.
FORBS							
<i>Achillea millefolium</i>	+ .12	+ .8	2.22	+ .8	+ .2	+ .6	+ .8
<i>Anemone patens</i>	.	.	.	.	.	.	+ .6
<i>Antennaria rosea</i>	*	+ .2	+ .2	*	+ .4	+ .6	+ .12
<i>Astragalus adsurgens</i>	.	.	.	+ .6	*	.	.
<i>Campanula rotundifolia</i>	+ .2	+ .4	1.16	1.18	+ .12	3.32	3.28
<i>Chenopodium album</i>	*	.	.	+ .2	.	.	.
<i>Fragaria virginiana</i>	+ .2	.	+ .2	.	.	.	.
<i>Galium boreale</i>	5.60	4.34	3.72	5.68	5.66	3.40	7.72
<i>Gentianella amarella</i>	*	+ .2	+ .4	.	.	.	1.14
<i>Geum triflorum</i>	.	.	.	.	.	+ .2	.
<i>Hackelia deflexa</i>	+ .6	.	.	.	.	.	.
<i>Heuchera richardsonii</i>	+ .4	*	*	*	+ .2	*	.
<i>Linum perenne</i>	.	.	.	.	.	.	.
<i>Melilotus officinalis</i>	.	.	.	.	.	.	.
<i>Mirabilis linearis</i>	.	.	.	+ .6	.	.	.
<i>Monarda fistulosa</i>	1.4	.	.	.	.	*	.
Mosses & Lichens	78.99	92.99	91.99	37.80	78.99	69.99	61.98
<i>Opuntia fragilis</i>	.	.	.	.	.	.	.
<i>Opuntia polyacantha</i>	.	.	.	.	.	.	.
<i>Oxytropis campestris</i>	.	.	+ .2	.	.	.	.
<i>Oxytropis lambertii</i>	.	1.4	.	.	.	.	+ .2
<i>Parietaria pennsylvanica</i>	.	.	.	.	*	.	.
Polypodiaceae	*	2.40	1.22	1.20	1.18	1.8	7.76
<i>Potentilla pensylvanica</i>	.	.	.	.	.	.	.
<i>Selaginella densa</i>	.	.	.	.	.	.	.
<i>Smilacina stellata</i>	2.18	+ .2	1.12	+ .4	1.10	*	1.10
<i>Solidago missouriensis</i>	+ .2	.	*	+ .6	.	.	1.6
<i>Taraxacum officinale</i>	.	.	+ .10	+ .10	+ .10	+ .6	+ .6
<i>Tragopogon dubius</i>	.	.	.	+ .2	.	.	.
<i>Vicia americana</i>	.	.	.	.	.	+ .2	.
<i>Viola adunca</i>	.	.	+ .6	1.20	+ .6	1.28	+ .14
<i>Viola</i> sp.	.	*	.	.	.	.	.
Species in microplots	20	15	19	19	15	17	20
Coverage of shrubs	59	6	30	9	12	4	2
Coverage of graminoids	59	62	68	73	59	83	80
Coverage of forbs	86	99	99	45	85	77	81
Total coverage	204	167	197	127	156	164	163



Table A9.—*Symphoricarpos occidentalis* community type

	Stand number				
	11	17	19	45	59
<b>Location</b>					
Quarter section	NW	NE	SE	SE	NW
Section	23	33	28	29	28
Township	140N	141N	141N	147N	148N
Range	102W	101W	101W	100W	100W
<b>Topographic position</b>					
Slope (%)	11	8	3	4	6
Aspect (°)	335	347	165	18	160
Elevation (m)	707	716	704	719	719
<b>Coverage/Frequency</b>					
<b>SHRUBS</b>					
<i>Artemisia cana</i>	.	+ .2	.	.	.
<i>Artemisia frigida</i>	.	+ .4	.	.	.
<i>Prunus americana</i>	.	.	1.2	.	.
<i>Prunus virginiana</i>	.	.	3.4	.	.
<i>Ribes odoratum</i>	6.8	.	.	.	.
<i>Rosa woodsii</i>	37.60	2.8	2.10	+ .2	.
<i>Symphoricarpos occidentalis</i>	89.99	85.99	95.99	98.99	98.99
<b>GRAMINOIDS</b>					
<i>Agropyron caninum</i>	.	1.6	11.22	1.4	.
<i>Agropyron smithii</i>	15.28	63.82	10.20	1.2	.
<i>Andropogon gerardii</i>	.	.	.	.	1.6
<i>Bromus inermis</i>	*	.	6.6	.	.
<i>Bromus tectorum</i>	.	.	.	+ .6	.
<i>Carex filifolia</i>	.	+ .2	.	.	.
<i>Elymus virginicus</i>	.	.	1.6	.	.
<i>Koeleria pyramidata</i>	.	+ .2	.	.	.
<i>Poa pratensis</i>	.	.	64.86	6.22	21.46
<i>Stipa viridula</i>	.	1.8	.	.	.
<b>FORBS</b>					
<i>Achillea millefolium</i>	+ .2	2.12	+ .6	.	+ .2
<i>Artemisia ludoviciana</i>	9.24	2.12	1.14	1.16	3.14
<i>Astragalus agrestis</i>	.	8.42	.	.	.
<i>Chenopodium album</i>	.	.	+ .2	+ .2	.
<i>Convolvulus arvensis</i>	1.4	.	+ .2	4.20	.
<i>Convolvulus sepium</i>	.	.	.	.	1.8
<i>Galium aparine</i>	+ .2	.	+ .2	+ .6	.
<i>Galium boreale</i>	.	.	6.16	8.24	10.42
<i>Hackelia deflexa</i>	.	.	.	.	+ .2
<i>Humulus lupulus</i>	.	.	9.14	.	.
<i>Lactuca oblongifolia</i>	11.44	1.22	38.90	+ .2	.
<i>Linum perenne</i>	.	2.18	.	.	.
<i>Linum rigidum</i>	+ .2	.	.	.	.
<i>Medicago sativa</i>	.	.	.	+ .2	.
<i>Parietaria pennsylvanica</i>	5.16	7.24	1.8	4.8	+ .4
<i>Ratibida columnifera</i>	+ .2	.	.	.	.
<i>Sisymbrium altissimum</i>	.	.	.	+ .2	.
<i>Taraxacum officinale</i>	+ .6	.	+ .2	+ .4	.
<i>Thalictrum venulosum</i>	.	.	.	.	+ .2
<i>Urtica dioica</i>	.	.	+ .2	.	.
Species in microplots	14	15	20	16	10
Coverage of shrubs	132	87	101	98	98
Coverage of graminoids	15	65	92	8	22
Coverage of forbs	26	22	55	17	14
Total coverage	173	174	248	123	134

## APPENDIX 2. TREE POPULATION ANALYSES

Population structure of trees in stands, by habitat types. Numbers of trees listed per 375 m<sup>2</sup>, with basal area for the stand as m<sup>2</sup>/ha given below the stand number. Abbreviations of tree species are as follows:

ACNE—*Acer negundo*  
CRCH—*Crataegus chrysocarpa*  
FRPE—*Fraxinus pennsylvanica*  
JUSC—*Juniperus scopulorum*  
PODE—*Populus deltoides*  
POTR—*Populus tremuloides*  
ULAM—*Ulmus americana*



Table A10.—Population structure of trees in stands, by habitat type

Stand and b.a.	Spp.	Diameter (at breast height) classes in dm							
		0-1		1-2	2-3	3-4	4-5	5-6	6-7
		<.5	>.5						
1. <i>Fraxinus pennsylvanica</i> / <i>Symphoricarpos occidentalis</i> habitat type									
29	FRPE	56							
39.5	JUSC	20	6	15					
	PODE						4	5	
30	FRPE	5							
23.3	JUSC	20	4	2	1				
	PODE						3	3	
51	FRPE	513	31	2					
63.1	JUSC	33	1	5					
	PODE							3	4
52	FRPE	489	8						
39.5	JUSC	6	9	4	1				
	PODE				4	3	3	1	
2. <i>Fraxinus pennsylvanica</i> / <i>Prunus virginiana</i> habitat type									
41	FRPE	332	20	28	5				
23.8	JUSC	6		1					
42	FRPE	306	11	12	8	1			
21.6	JUSC			1					
57	FRPE	15	5	17	9	4			
35.9	ULAM	66			2	1			
	JUSC	2							
58	FRPE	5	1	15	3				
25.1	ULAM	21	8	15	4				
	ACNE	8	1	1					
	JUSC	2							
3. <i>Populus tremuloides</i> / <i>Betula occidentalis</i> habitat type									
43	POTR	81	58	9					
14.7	FRPE	47	10	2					
	JUSC	7							
60	POTR	82	35	28	1				
21.1	FRPE	41							
	ULAM	9	1	2					
	CRCH	1							
67	POTR	136	15	32					
20.9	FRPE	61	6	2					
	JUSC	1							
68	POTR	22	2	17	22				
41.8	FRPE	92	5	6					
	ACNE	13	5						
4. <i>Juniperus scopulorum</i> / <i>Oryzopsis micrantha</i> habitat type									
33	JUSC	47	21	20	1	3			
29.6	FRPE	62	15	5					
34	JUSC	126	60	30	1				
23.4									
37	JUSC	112	82	24					
22.1	FRPE	17	1						
38	JUSC	98	86	30					
25.4	FRPE	6							
49	JUSC	207	108	10	2				
22.2	FRPE	1							
50	JUSC	158	43	26	4				
22.7	FRPE	2							
53	JUSC	232	59	21	3				
22.4	FRPE	10							

### APPENDIX 3. SOIL ANALYSES

Table A11.—Results of soil analyses of the upper 1 dm of mineral soil from each stand. Data are arranged according to habitat types

Stand	pH	C.E.C. (meq/ 100 g)	Free lime <sup>1</sup>	Extractable cations (meq/100 g)			P (ppm)	N (%)	O.M. <sup>2</sup> (%)	Mechanical analysis (percent)			Texture
				Ca	Mg	Na				Sand	Silt	Clay	
Stipa comata/Carex filifolia habitat type													
27	6.2	-	-	-	-	-	-	0.174	2.6	56.4	34.0	9.6	Sandy loam
35	6.9	-	-	-	-	-	-	0.116	1.8	64.4	29.6	6.0	Sandy loam
36	6.7	6.84	-	7.65	3.58	0.14	4.5	0.097	2.0	71.4	22.6	6.0	Sandy loam
39	7.5	6.47	+	16.67	3.54	0.18	3.0	0.107	1.4	74.0	18.0	8.0	Sandy loam
40	7.1	-	-	-	-	-	-	0.073	1.0	79.0	13.0	8.0	Loamy sand
56	7.1	-	-	-	-	-	-	0.114	1.6	53.6	35.6	10.8	Sandy loam
61	6.8	9.61	+	51.10	1.73	0.07	4.0	0.081	1.3	86.0	7.6	6.4	Loamy sand
Agropyron smithii/Carex filifolia habitat type													
3	7.7	-	-	-	-	-	-	0.159	2.4	32.4	45.0	22.6	Loam
5	7.7	-	-	-	-	-	-	0.180	2.6	39.4	40.0	20.6	Loam
14	7.6	18.38	+++	39.05	2.47	0.12	0.5	0.178	2.7	38.8	37.6	23.6	Loam
18	7.7	13.83	+++	26.66	3.98	0.10	0.5	0.178	2.4	45.2	35.2	19.6	Loam
20	7.7	-	-	-	-	-	-	0.171	2.2	50.4	32.0	17.6	Loam
21	7.7	-	-	-	-	-	-	0.149	2.0	43.4	40.6	16.0	Loam
48	7.6	-	-	-	-	-	-	0.153	1.8	50.0	36.0	14.0	Loam
54	6.7	28.47	-	11.54	4.79	0.11	1.0	0.181	2.1	44.0	41.0	15.0	Loam
62	7.6	-	-	-	-	-	-	0.137	2.0	35.0	40.6	24.4	Loam
Andropogon scoparius/Carex filifolia habitat type													
8	7.7	-	-	-	-	-	-	0.164	2.6	66.4	26.0	7.6	Sandy loam
15	7.7	9.07	+++	26.05	1.35	0.11	0.5	0.158	2.1	61.2	31.2	7.6	Sandy loam
22	7.7	-	-	-	-	-	-	0.237	3.0	34.4	49.6	16.0	Loam
23	7.7	14.11	++	30.01	3.12	0.15	3.0	0.190	2.4	48.4	41.6	10.0	Loam
24	7.7	-	-	-	-	-	-	0.155	2.2	52.4	40.0	7.6	Sandy loam
46	7.7	-	-	-	-	-	-	0.154	2.1	22.6	55.0	22.4	Silt loam
55	7.8	-	-	-	-	-	-	0.087	1.1	23.6	49.6	26.8	Loam
63	7.6	-	-	-	-	-	-	0.142	2.0	57.0	28.6	14.4	Loam
64	7.6	11.32	+++	24.45	3.51	0.07	0.5	0.123	1.8	48.4	38.6	13.0	Loam
Juniperus horizontalis/Andropogon scoparius habitat type													
9	7.6	-	-	-	-	-	-	0.112	2.4	40.4	37.4	22.2	Loam
10	7.6	17.88	+++	54.16	3.30	0.25	0.5	0.232	3.4	26.4	44.0	29.6	Clay loam
12	7.4	26.67	++	34.03	2.74	0.04	1.0	0.219	3.0	48.8	29.6	21.6	Loam
44	7.3	39.13	+	36.89	6.37	0.27	2.0	0.274	4.2	48.0	41.0	11.0	Loam
47	7.5	-	-	-	-	-	-	0.138	2.5	37.6	34.0	28.4	Clay loam
Artemisia tridentata/Agropyron smithii habitat type													
4	7.4	26.42	+	43.17	6.97	1.57	6.0	0.147	3.6	23.0	35.4	41.6	Clay
13	7.1	-	-	-	-	-	-	0.146	3.5	24.8	71.6	3.6	Silt loam
16	7.5	10.31	+	20.45	5.81	0.71	2.0	0.165	2.2	39.2	35.2	25.6	Loam
28	7.6	31.02	+++	32.04	4.61	1.30	1.0	0.177	2.5	26.0	41.6	32.4	Clay loam
Artemisia cana/Agropyron smithii habitat type													
1	7.8	15.98	+++	39.22	2.69	2.61	2.5	0.100	1.8	18.4	50.0	31.6	Silty clay loam
6	7.8	-	-	-	-	-	-	0.157	2.0	14.4	50.0	35.6	Silty clay loam
7	7.8	-	-	-	-	-	-	0.137	1.9	13.4	52.0	34.6	Silty clay loam
26	7.7	-	-	-	-	-	-	0.098	2.9	40.4	46.6	13.0	Loam
66	8.0	-	-	-	-	-	-	0.069	1.7	17.8	63.6	18.6	Silt loam
69	7.8	11.31	+++	23.61	3.21	1.11	24.0	0.071	1.4	13.8	61.6	24.6	Silt loam
70	7.7	-	-	-	-	-	-	0.046	1.7	58.8	29.6	11.6	Sandy loam
2	7.6	11.22	+++	33.52	2.42	0.11	1.0	0.101	2.1	38.4	29.0	32.6	Clay loam
25	7.6	-	-	-	-	-	-	0.098	2.2	34.4	47.6	18.0	Loam
31	7.7	8.82	+++	20.18	3.39	0.25	0.5	0.082	2.0	38.0	42.0	20.0	Loam
32	7.7	-	-	-	-	-	-	0.055	1.6	36.0	54.0	10.0	Silt loam
65	7.6	-	-	-	-	-	-	0.088	3.6	34.8	50.6	14.6	Silt loam
Fraxinus pennsylvanica/Symphoricarpos occidentalis habitat type													
29	7.6	17.57	++	21.13	3.13	0.31	2.5	0.138	2.6	31.0	51.0	18.0	Silt loam
30	7.6	-	-	-	-	-	-	0.089	1.7	44.0	41.0	15.0	Loam
51	7.5	28.21	++	30.31	6.42	0.59	1.0	0.152	2.6	31.0	35.0	34.0	Clay loam
52	7.5	17.14	+	27.91	6.61	0.23	2.0	0.155	3.4	22.0	50.6	27.4	Clay loam



Table A11.—Results of soil analyses of the upper 1 dm of mineral soil from each stand.  
Data are arranged according to habitat types—Continued

Stand	pH	C.E.C. (meq/ 100 g)	Free lime <sup>1</sup>	Extractable cations (meq/100 g)			P (ppm)	N (%)	O.M. <sup>2</sup> (%)	Mechanical analysis (percent)			Texture
				CA	Mg	Na				Sand	Silt	Clay	
Fraxinus pennsylvanica/Prunus virginiana habitat type													
41	7.6	—	—	—	—	—	—	0.202	2.9	33.0	33.0	34.0	Clay loam
42	7.6	15.25	+ + +	32.23	4.97	0.25	0.5	0.154	2.5	15.0	39.0	46.0	Clay
57	6.0	25.53	—	18.52	6.83	0.12	9.5	0.362	4.1	34.6	45.6	19.8	Loam
58	6.4	24.83	—	17.72	3.89	0.10	9.5	0.406	3.6	38.0	43.2	18.8	Loam
Populus tremuloides/Betula occidentalis habitat type													
43	7.2	12.94	—	13.56	4.89	0.13	3.0	0.099	2.3	35.0	39.0	26.0	Loam
60	5.7	—	—	—	—	—	—	0.221	3.3	48.0	34.6	17.4	Loam
67	6.0	18.23	—	13.98	6.05	0.08	9.0	0.111	1.9	64.4	24.0	11.6	Sandy loam
68	7.1	16.66	—	12.53	4.06	0.11	1.0	0.103	1.6	32.8	39.6	27.6	Clay loam
Juniperus scopulorum/Oryzopsis micrantha habitat type													
33	6.7	—	—	—	—	—	—	0.295	4.3	57.0	33.0	10.0	Sandy loam
34	7.9	17.58	+ + +	27.62	8.10	1.57	0.5	0.130	2.3	25.0	39.0	36.0	Clay loam
37	7.8	—	—	—	—	—	—	0.167	3.0	27.0	34.0	39.0	Clay loam
38	7.6	30.91	—	19.43	13.91	3.55	0.5	0.203	3.3	31.0	30.0	39.0	Clay loam
49	7.6	—	—	—	—	—	—	0.180	3.0	26.0	39.0	35.0	Clay loam
50	7.8	—	—	—	—	—	—	0.149	3.3	35.0	36.0	29.0	Clay loam
53	7.8	23.09	+	25.80	8.22	1.32	0.5	0.164	2.6	44.0	32.0	24.0	Loam
Symphoricarpos occidentalis community type													
11	7.4	—	—	—	—	—	—	0.259	3.1	45.8	28.6	25.6	Loam
17	7.5	15.93	+ + +	28.11	2.90	0.17	2.5	0.222	3.0	47.2	31.2	21.6	Loam
19	7.5	—	—	—	—	—	—	0.341	4.2	18.0	48.0	34.0	Silty clay loam
45	6.2	23.90	—	16.43	5.92	0.10	2.5	0.286	3.4	32.4	45.6	22.0	Loam
59	5.8	—	—	—	—	—	—	0.273	3.3	35.0	46.2	18.8	Loam

<sup>1</sup>Free lime determined subjectively. + = low, ++ = medium, +++ = high.

<sup>2</sup>Organic matter.

Table A12.—Gravel content (percent by weight) of soil samples taken from the upper 1 dm of mineral soil from each stand

Habitat type	Stand	Gravel content
<i>Stipa comata</i> / <i>Carex filifolia</i>	61	11.4
<i>Andropogon scoparius</i> / <i>Carex filifolia</i>	22	6.7
<i>Andropogon scoparius</i> / <i>Carex filifolia</i>	23	36.2
<i>Andropogon scoparius</i> / <i>Carex filifolia</i>	24	30.3
<i>Juniperus horizontalis</i> / <i>Andropogon scoparius</i>	10	6.5
<i>Juniperus horizontalis</i> / <i>Andropogon scoparius</i>	44	29.2
<i>Artemisia tridentata</i> / <i>Agropyron smithii</i>	28	4.6
<i>Artemisia cana</i> / <i>Agropyron smithii</i>	2	16.0





Hansen, Paul L., George R. Hoffman, and Ardell J. Bjugstad. 1984. The vegetation of Theodore Roosevelt National Park, North Dakota: A habitat type classification. General Technical Report RM-113, 35 p. Rocky Mountain Forest and Range Experiment Station, Fort Collins, Colo.

Vegetation of the Theodore Roosevelt National Park, North Dakota, was characterized according to habitat type based on concepts and methods developed by Daubenmire. Ten habitat types—three grassland and seven wooded types—were described including a key to identify the habitat types.

**Keywords:** Vegetation classification, habitat type, *Stipa comata*, *Carex filifolia*, *Andropogon scoparius*, *Juniperus horizontalis*, *Artemisia tridentata*, *Agropyron smithii*, *Artemisia cana*, *Fraxinus pennsylvanica*, *Symphoricarpos occidentalis*, *Prunus virginiana*, *Populus tremuloides*, *Betula occidentalis*, *Juniperus scopulorum*, *Oryzopsis micrantha*

Hansen, Paul L., George R. Hoffman, and Ardell J. Bjugstad. 1984. The vegetation of Theodore Roosevelt National Park, North Dakota: A habitat type classification. General Technical Report RM-113, 35 p. Rocky Mountain Forest and Range Experiment Station, Fort Collins, Colo.

Vegetation of the Theodore Roosevelt National Park, North Dakota, was characterized according to habitat type based on concepts and methods developed by Daubenmire. Ten habitat types—three grassland and seven wooded types—were described including a key to identify the habitat types.

**Keywords:** Vegetation classification, habitat type, *Stipa comata*, *Carex filifolia*, *Andropogon scoparius*, *Juniperus horizontalis*, *Artemisia tridentata*, *Agropyron smithii*, *Artemisia cana*, *Fraxinus pennsylvanica*, *Symphoricarpos occidentalis*, *Prunus virginiana*, *Populus tremuloides*, *Betula occidentalis*, *Juniperus scopulorum*, *Oryzopsis micrantha*

Hansen, Paul L., George R. Hoffman, and Ardell J. Bjugstad. 1984. The vegetation of Theodore Roosevelt National Park, North Dakota: A habitat type classification. General Technical Report RM-113, 35 p. Rocky Mountain Forest and Range Experiment Station, Fort Collins, Colo.

Vegetation of the Theodore Roosevelt National Park, North Dakota, was characterized according to habitat type based on concepts and methods developed by Daubenmire. Ten habitat types—three grassland and seven wooded types—were described including a key to identify the habitat types.

**Keywords:** Vegetation classification, habitat type, *Stipa comata*, *Carex filifolia*, *Andropogon scoparius*, *Juniperus horizontalis*, *Artemisia tridentata*, *Agropyron smithii*, *Artemisia cana*, *Fraxinus pennsylvanica*, *Symphoricarpos occidentalis*, *Prunus virginiana*, *Populus tremuloides*, *Betula occidentalis*, *Juniperus scopulorum*, *Oryzopsis micrantha*

Hansen, Paul L., George R. Hoffman, and Ardell J. Bjugstad. 1984. The vegetation of Theodore Roosevelt National Park, North Dakota: A habitat type classification. General Technical Report RM-113, 35 p. Rocky Mountain Forest and Range Experiment Station, Fort Collins, Colo.

Vegetation of the Theodore Roosevelt National Park, North Dakota, was characterized according to habitat type based on concepts and methods developed by Daubenmire. Ten habitat types—three grassland and seven wooded types—were described including a key to identify the habitat types.

**Keywords:** Vegetation classification, habitat type, *Stipa comata*, *Carex filifolia*, *Andropogon scoparius*, *Juniperus horizontalis*, *Artemisia tridentata*, *Agropyron smithii*, *Artemisia cana*, *Fraxinus pennsylvanica*, *Symphoricarpos occidentalis*, *Prunus virginiana*, *Populus tremuloides*, *Betula occidentalis*, *Juniperus scopulorum*, *Oryzopsis micrantha*









Rocky  
Mountains



Southwest



Great  
Plains

U.S. Department of Agriculture  
Forest Service

## Rocky Mountain Forest and Range Experiment Station

The Rocky Mountain Station is one of eight regional experiment stations, plus the Forest Products Laboratory and the Washington Office Staff, that make up the Forest Service research organization.

### RESEARCH FOCUS

Research programs at the Rocky Mountain Station are coordinated with area universities and with other institutions. Many studies are conducted on a cooperative basis to accelerate solutions to problems involving range, water, wildlife and fish habitat, human and community development, timber, recreation, protection, and multiresource evaluation.

### RESEARCH LOCATIONS

Research Work Units of the Rocky Mountain Station are operated in cooperation with universities in the following cities:

Albuquerque, New Mexico  
Flagstaff, Arizona  
Fort Collins, Colorado\*  
Laramie, Wyoming  
Lincoln, Nebraska  
Rapid City, South Dakota  
Tempe, Arizona

\*Station Headquarters: 240 W. Prospect St., Fort Collins, CO 80526